

## Properties, various functions and application of bacterial melanins – some properties and possible application of melanin produced by *Azotobacter chroococcum*

Monika Koziel

Department of Agricultural Microbiology, Institute of Soil Science and Plant Cultivation – State Research Institute  
ul. Czartoryskich 8, 24-100 Puławy, POLAND

**Abstract.** Melanins are a group of macromolecular pigments formed as a result of oxidative polymerization of phenolic and indole compounds. They are considered the most common, heterogeneous, resistant and evolutionarily oldest pigments found in nature. Melanins occurred very early in the evolution of various groups of organisms. They have already been found in dinosaur, bird and primitive cephalopod fossils. Today, melanins are found in every kingdom of living organisms and play an important role in the processes of reproduction, thermoregulation, chemoprotection and camouflage. In addition to the important functions they perform in organisms, they exhibit a different chemical structure and are characterized by a wide range of colors, from black-brown to yellow-red. Differences in chemical structure have become the criterion for dividing melanins into four groups, i.e. eumelanin, allomelanin, pheomelanin and neuromelanin. Production of melanins is characteristic for many microorganisms, including free-living *Azotobacter* bacteria. The genus *Azotobacter* comprises eight species and only *Azotobacter bryophylli* does not produce pigments. *Azotobacter chroococcum*, the most abundant in soils all over the world, produces a dark brown melanin pigment non-diffusible into the substrate. Melanins synthesized by this species of bacteria increased the growth of some plants and detoxification of soils and waters polluted with heavy metals. In addition, the method of obtaining melanin produced by *A. chroococcum* is simple and relatively cheap compared to the cost of obtaining synthetic melanins, which gives the opportunity to conduct further research on the use of this pigment in biotechnology and molecular biology. In work it was describe the physicochemical properties, various functions and possible applications of bacterial melanins in various industries. The publication also summarizes the current knowledge on some properties and the possibility of use in bioremediation of soils and waters contaminated with heavy metals melanins synthesized by *Azotobacter chroococcum*.

**Key words:** bacterial melanin, polyphenol oxidase, stress protection, biosynthesis, *Azotobacter chroococcum*

Corresponding author:

Monika Koziel  
e-mail: mkoziel@iung.pulawy.pl  
phone: +48 81 4786 952

### INTRODUCTION

Natural pigments are extracted from biological material of plant, animal and microbial origin (Stolarzewicz et al., 2012). Many bacterial species can synthesize pigments. Colorful bacteria outcompete species and mutants that do not produce pigments. In the case of bacteria, the benefits of pigment production are the acquisition of energy and food, the extraction of ferric from the habitat, protection from the harmful effects of high-concentration visible light, ultraviolet radiation, extreme temperatures and compounds with antibacterial effects. It is noteworthy that some pigments have antibiotic activity as well as virulence factors for pathogenic bacteria. The pigmentation of microorganisms is essential in clinical diagnostics, as it facilitates species identification (Wolska et al., 2010).

Melanins are a group of macromolecular, black or brown in color pigments formed as a result of a multi-step oxidative polymerization of phenolic and indole compounds. The most common substrate for the biosynthesis of these compounds is the amino acid tyrosine, dihydroxyindole, dihydroxynaphthalene and catechol. Melanins exhibit specific physical and chemical properties. They reduce many compounds, including silver ions, ferrocyanide, and potassium permanganate, have UV absorption and scattering abilities, exhibit antioxidant properties, and are excellent chelators of metal cations (Saud, Alaubydi, 2016). Production of melanins is characteristic for many microorganisms, mainly soil bacteria, some parasitic bacteria and fungi, and marine algae. Among soil bacteria, melanins are produced by actinomycetes (some species of *Streptomyces* (Paget et al., 1994)), bacterial species belonging to the genus *Azotobacter* (Shivprasad, Page, 1989; Aquilanti et al., 2004a), *Azospirillum* (Sadasivan, Neyra, 1987) and some strains of *Rhizobium* (Hynes et al., 1988). Among marine algae, *Shewanella* (Turick et al., 2008), *Shewanella colwelliana* (Kotob et al., 1995), and *Marinomonas mediterranea* (Lopez-Serrano et al., 2004) possess this ability. On the other hand, among pathogenic bacteria, melanin is

most often produced by species, i.e., *Burkholderia cepacia* (Zughaier et al., 1999), *Klebsiella pneumoniae* (Hawkins, Johnston, 1988), *Legionella pneumophila* (Baine et al., 1978), *Mycobacterium leprae* (Prabhakaran et al., 1968), *Proteus mirabilis* (Agodi et al., 1996) and *Vibrio cholerae* (Coyne, Al-Harhi, 1992).

Bacteria belonging to the genus *Azotobacter* are free-living, strict aerobes that require oxygen. Of the species of this genus, *Azotobacter chroococcum* is the most widely distributed in soils worldwide, and it probably also dominates soils in Poland (Döbereiner, 1995; Martyniuk, Martyniuk, 2003; Lenart, 2012). Colonies of these bacteria are large, convex with an irregular shape, mucilaginous, and darkening after several days of culture, which is associated with the production of a dark brown melanin pigment non-diffusible into the substrate. The aspects of melanin production by *Azotobacter chroococcum* have been described in quite some detail in the literature (Thompson, Skerman, 1979; Shivprasad, Page, 1989; Gospodaryov, Lushchak, 2011; Banerjee et al., 2014).

The paper systematizes current reports on the conditions of melanin production by bacteria, including by *Azotobacter chroococcum*, some of its physicochemical properties and possible applications in biotechnology. The work presents, in a synthetic form, the current state of knowledge in this area.

#### PHYSICOCHEMICAL PROPERTIES OF MELANINS

Melanins are a diverse group of pigments synthesized in living prokaryotic and eukaryotic organisms by hydroxylation and polymerization of organic compounds. Melanin production is one of the universal features of the adaptation of living organisms to changing environmental conditions. The presence of different types of melanin in representatives of almost every major taxon indicates the evolutionary significance of melanogenesis (Plonka, Grabacka, 2006).

All the pigments belonging to the melanin group are characterized by their high molecular weight, irregular and three-dimensional amorphous structure and negative charge. There is no single, strict definition of melanins due to the high heterogeneity of these compounds (color, molecular weight, composition, origin and function). The most commonly used definition is: „heterogeneous polymers formed by the oxidation of phenolic compounds and further polymerization of intermediate compounds and resulting quinones” (Solano, 2014). The diverse and heterogeneous structure of melanins is due to the ubiquitous sources of their origin. In addition, melanin’s physicochemical properties make it even more difficult to accurately identify and characterize its structure (Pralea et al., 2019). Among the characteristic qualities of melanins that allow them to be distinguished from other compounds are insolubility in most common solvents, loss of color under oxidizing agents, resistance to degradation under cold and

hot acids, ability to directly reduce ammoniacal AgNO<sub>3</sub> solution, solubility in alkali solutions, positive reaction to polyphenols (Piattelli et al., 1965; Nicolaus, 1968; Nosanchuk et al., 2015; Pralea et al., 2019). A dark color usually characterizes melanins, but the palette ranges from blackish brown to yellowish red. This variation in color and hue is due to different abilities to absorb and scatter light. According to literature data, the smaller the melanin granules, the lighter their coloration (Prota, 1992).

#### TYPES OF MELANINS

The melanin-producing ability is widespread among microorganisms. From a chemical point of view, the common feature of microbial melanins is that they are the product of oxidative polymerization of various phenolic substances (Przemyslaw, May, 2006). Melanins are heterogeneous polymers of dihydroxyindole (DHI) and dihydroxyindolecarboxylic acid (DHICA) monomers linked by heterogeneous non-hydrolyzing bonds (Crippa et al., 1989; Cheng et al., 1994). Differences in chemical structure have become the criterion for dividing melanins into four groups (Table 1), i.e.:

- eumelanins – blackish brown insoluble pigments formed by the oxidation of tyrosine (and/or phenylalanine) to 3,4-dihydroxyphenylalanine (DOPA) and dopaquinone, which are further converted to 5,6-dihydroxyindole or 5,6-dihydroxyindole-2-carboxylic acid. Eumelanins are most common in animals and humans (Langfelder et al., 2003; Plonka, Grabacka, 2006),
- allomelanins – are the least studied and the most heterogeneous group of polymers formed by the oxidation of di-hydroxynaphthalene or tetrahydroxynaphthalene to phaeomelanin,  $\gamma$ -glutamine-4-hydroxybenzene, catechol and 4-hydroxyphenylacetic acid. Plants, fungi and microorganisms produce allomelanins. This group includes the following subgroups of melanins: catechol melanin (in plants), DHN melanin and phaeomelanin (in bacteria and fungi) (Nicolaus et al., 1964; Funa et al., 1999; Jacobson, 2000; Plonka, Grabacka, 2006; Cordero, Casadevall, 2017),
- pheomelanins – yellow-red pigments soluble in alkaline solutions. This variety of melanins is formed by biochemical transformations involving a change in eumelanin synthesis. The onset of synthesis of both compounds is similar; however, in the case of pheomelanin production, 3,4-dihydroxyphenylalanine undergoes cysteinylolation, either directly or via glutathione. Cysteinyl-3,4-dihydroxyphenylalanine, the final product of the reaction, further polymerizes into various benzothiazine derivatives. This is a type of animal melanin found in red hair, freckles or feathers (Nappi, Ottaviani, 2000; Plonka, Grabacka, 2006),
- neuromelanins – dark pigments synthesized in human neurons by oxidation of dopamine and other catecholamine precursors (Fedorov et al., 2005).

Table 1. Melanins – types, source of origin and precursors (Tran-Ly et al., 2020).

Type of melanin	Source of origin	Precursor
Eumelanin	animals, bacteria, fungi	tyrosine, 3,4-dihydroxy-L-phenylalanine (L-DOPA)
Pheomelanin	animals	5-S-cysteinyl-DOPA
Neuromelanin	humans (brain)	dopamine, 5-S-cysteinyl-dopamine
Catechol melanin	plants	catechol
DHNmelanin	bacteria, fungi	1,8-dihydroxynaphtalene (DHN)
Phaeomelanin	bacteria, fungi	homogentisinic acid

#### ECOLOGICAL AND PHYSIOLOGICAL ROLE OF BACTERIAL MELANINS

The capability of microorganisms to produce pigments is a trait acquired in the course of evolution, increasing their survival opportunities in the external environment. The pigments are designed to protect microorganisms from the adverse effects of various physicochemical factors (Liu, Nizet, 2009). Due to the multifunctional nature of melanins, they are used as a) antioxidants and radical scavengers (Keith et al., 2007; Ju et al., 2011; Le Na et al., 2019), b) photo protectors that effectively absorb and dissipate solar radiation in the form of heat (d'Ischia et al., 2015), (c) absorbers that chelate metals and bind organic compounds (Banerjee et al., 2014; Karlsson, Lindquist, 2016; Tran-Ly et al., 2020a), and (d) organic semiconductors (Bothma et al., 2008). Besides, these pigments protect cells from hydrolytic enzymes and the enzymes themselves from proteases (Valeru et al., 2009; Saud, Alaubydi, 2016). In addition to performing the earlier functions, melanin is considered a biocompatible and environmentally friendly compound, as it is naturally synthesized by most living organisms (Tran-Ly et al., 2020b). The link between melanin synthesis and increased virulence of pathogenic bacteria has been widely reported in the literature (Nosanchuk, Casadevall, 2003; Plonka, Grabacka, 2006). The pigment contributes to the virulence of microorganisms by decreasing the susceptibility of pathogens to host defense mechanisms and influences the host immune response (Nosanchuk, Casadevall, 2006). Melanins synthesized by water-dwelling *Vibrio cholerae* bacteria protect cells from high temperature and osmotic stress. The protective role of melanins under osmotic stress is related to their ability to absorb K<sup>+</sup> and Na<sup>+</sup> cations, which prevents cell dehydration (Coyne, Al-Harthi., 1992). A study by Valeru et al. (2009) shows that the color mutant of *Vibrio cholerae* has enhanced infectivity and increased resistance to UV radiation due to the absorption of the energy of this radiation by melanin. On the other hand, Patel et al. (1996) showed that the melanin of *Bacillus thuringiensis* increases the toxicity of this bacterium to insects while protecting it from the adverse effects of UV radiation. In addition, resistance *B. thuringiensis* to UV radiation has been linked to the presence of melanin in the sheaths of endospores, while adsorption or incorporation of melanin into crystals of the toxic protein formed during sporulation increases the stability of the insecticide.

In turn, the ability of melanins to bind heavy metals is associated with numerous carboxyl, phenolic, hydroxyl and amino groups in their structure, which are sites of binding or adsorption for metal ions. This can reduce the effectiveness of antifungal and antibacterial agents containing metal ions in eliminating melanin-producing microorganisms (Patel et al., 1996).

The literature has reported the potent antioxidant properties of melanins and their ability to stabilize free radicals and bind unpaired electrons to increase the virulence of colored strains compared to non-colored strains. The above relation was observed in some colored strains of *Burkholderia cenocepacia* (Keith et al., 2007) and the plant pathogen *Ralstonia solanacearum* (Ahmad et al.,

2016). Piñero et al. (2007) found that melanin in the symbiotic bacterium *Rhizobium etli* plays a key role in the first stages of wart formation when the bacteria must deal with reactive oxygen species and phenolic compounds produced by plants. These dyes are also involved in interactions between bacterial biofilms and other organisms. Melanin secreted by *Pseudoalteromonas lipolytica* biofilms prevents their colonization by clam larvae (Zeng et al., 2017) and in the case of a *Vibrio cholerae* biofilm increases the production of reactive oxygen species, thereby protecting the microorganism from predation by the amoeba species *Acanthamoeba castellanii* (Noorian et al., 2017).

Melanins also acquire iron from the environment (Gospodaryov, Lushchak, 2011). Most bacteria obtain this element through the production of siderophores. Chatfield and Cianciotto (2007) described a system of ferric procurement by a bacterium of the *Legionella pneumophila* species with the participation of pyomelanin, associated with ferric reductase activity. Melanins produced by bacteria belonging to the genera: *Azotobacter*, *Burkholderia*, *Pseudomonas*, *Klebsiella*, *Serratia* and *Vibrio* are also involved in iron acquisition. The mechanism of melanin-mediated ferric oxide reduction has been described in the Gram-negative marine bacteria *Shewanella alge* and the fungus *Cryptococcus neoformans*. Among pathogenic bacteria, *Pseudomonas aeruginosa* can reduce ferric extracellularly with pyomelanin (Chatfield, Cianciotto, 2007). Some microorganisms, as a result of melanin synthesis, are able to survive in extreme environments by adapting to rapidly changing conditions. Examples of such microorganisms include *Streptomyces cyaneofuscatus* isolated from the desert soils of Algeria (Harir et al., 2018), *Bacillus weihenstephanensis* from the soils of northeastern Poland (Drewnowska et al., 2015), *Lysobacter oligotrophicus* obtained from Antarctic sites (Kimu-

ra et al., 2015) and *Aeromonas salmonicida* subsp. *pectinolytica* from heavily polluted waters of the Matanza River in Argentina (Pavan et al., 2000).

SYSTEMATIC AFFILIATION AND PIGMENT  
PRODUCTION BY SPECIES OF THE GENUS  
*AZOTOBACTER*

The genus *Azotobacter* was identified by Beijerinck in 1901. Bacteria of the genus *Azotobacter* belongs to the family *Pseudomonadaceae*, included in the subclass  $\gamma$ -Proteobacteria (Tchan, New, 1984; Özen, Ussery, 2012; Rubio et al., 2013; Robson et al., 2015). Currently, 8 species and 4 subspecies are known within the genus *Azotobacter* (De Smedth et al., 1980; Tchan, New, 1984; Aquilanti et al., 2004b; Jin et al., 2020).

- *Azotobacter armeniacus* (Thompson, Skerman, 1979),
- *Azotobacter beijerinckii* (Lipman, 1904),
- *Azotobacter bryophylli* (Liu et al., 2019),
- *Azotobacter chroococcum* (Beijerinck, 1901),
  - *Azotobacter chroococcum* subsp. *chroococcum* (Jin et al., 2020),
  - *Azotobacter chroococcum* subsp. *isscasi* (Jin et al., 2020),
- *Azotobacter nigricans* (Krasilnikov, 1949),
  - *Azotobacter nigricans* subsp. *achromogenes* (Thompson, Skerman, 1979),
  - *Azotobacter nigricans* subsp. *nigricans* (Howey et al., 1990),
- *Azotobacter paspali* (Döbereiner, 1966),
- *Azotobacter salinestrus* (Page, Shivprasad, 1991),
- *Azotobacter vinelandii* (Lipman, 1903).

Of the species mentioned above, *Azotobacter bryophylli* is a colorless species, while the other species show the ability to produce melanin pigments of different colors (Table 2).

Table 2. Pigments produced by species belonging to the genus *Azotobacter* (Tchan, New, 1984; Page, Shivprasad, 1991; Aquilanti et al., 2004a; Liu et al., 2019).

Species	Pigment
<i>A. armeniacus</i>	brown-black pigment diffusing into the substrate
<i>A. beijerinckii</i>	yellow to light brown pigment not diffusing into the substrate
<i>A. bryophylli</i>	does not produce any pigment
<i>A. chroococcum</i>	dark brown melanin pigment not diffusing to the substrate
<i>A. nigricans</i>	brownish-black to reddish-purple pigment diffusing into the substrate
<i>A. paspali</i>	yellow fluorescent pigment diffusing into the substrate
<i>A. salinestrus</i>	dark brown pigment not diffusing into substrate
<i>A. vinelandii</i>	yellow-green fluorescent pigment diffusing to substrate

PROPERTIES OF MELANIN SYNTHESIZED  
BY AZOTOBACTER CHROOCOCCUM AND POSSIBLE APPLICATIONS.

Free-living  $N_2$  assimilators of the genus *Azotobacter* have been the subject of much research for decades. They have become model microorganisms in studies of the biochemistry and energetics of  $N_2$  fixation, as well as in studies of the spatial structure and function of nitrogenase and the genetic regulation of biological, atmospheric nitrogen fixation (Paul, Clark, 2000).

The ability of these bacteria to produce melanins is widely reported in the literature (Tchan, New, 1984; Shivprasad, Page, 1989; Gospodaryov, Lushchak, 2011; Banerjee et al., 2014). The compound used by *Azotobacter chroococcum* to

produce melanin is catechol. The most significant amounts of this pigment are produced under aerobic conditions because oxygen is one of the substrates of the enzyme polyphenol oxidase (Shivprasad, Page, 1989). Colonies of *A. chroococcum* bacteria darken after several days of culture due to the accumulation of dark brown melanin pigment that does not diffuse into the medium (Tchan, New, 1984; Aquilanti et al., 2004a) (Fig. 1). Melanin produced by this bacterial species belongs to the allomelanins group. Its synthesis does not require the presence of precursor molecules (e.g., tyrosine, dihydroxyphenylalanine, catechol) in the culture medium, as the bacteria produce these as the cells age (Bortels, Henkel, 1968). Melanin is produced after about five days of culture, staining most liquid medium or forming dark halos around bacterial colonies on solid medium. Melanin formation has been studied in several strains of *A. chroococcum* (Shivprasad, Page, 1989; Herter et al., 2011; Banerjee et al., 2014). Gospodaryov, Lushchak (2011) observed increased melanin production on the Ashby substrate with benzoic acid as a carbon source.

On the other hand, Shivprasad, Page (1989) noticed an inhibitory effect of benzoic acid and a stimulating effect of copper and iron ions on melanin production by *Azotobacter chroococcum*. They found that pigment synthesis was significantly enhanced at specific concentrations of copper sulfate in the medium. This is confirmed by a study by Gospodaryov, Lushchak (2011), who noted that copper sulfate at a concentration of 10  $\mu$ M added to the substrate increased melanin production, while higher concentrations inhibited the process. Excessively high concentrations of copper sulfate can inhibit bacterial growth and reduce the efficiency of this pigment syn-




Figure 1. 5-day-old colonies of the reference strain *Azotobacter chroococcum* DSM 281.

thesis. The studies discussed above show that melanogenesis in *Azotobacter chroococcum* is catalyzed by a copper-dependent polyphenol oxidase.

Like all melanins, melanin produced by *A. chroococcum* is soluble in alkaline reagents (Shivprasad, Page, 1989; Lin et al., 2005). Shivprasad and Page (1989) identified the dye extracted from *A. chroococcum* cells as melanin based on the following characteristics: solubility in hot 0.5 M NaOH and 1 M Na<sub>2</sub>CO<sub>3</sub>, and insolubility in cold water, hot water, ethanol, chloroform, acetone and cold 0.5 M NaOH. Melanin synthesized by this species can bind calcium ions. Alkalinization of the medium containing calcium ions results in the formation of black floes that are a melanin complex with Ca(OH)<sub>2</sub>. The ability to bind calcium is not a distinguishing feature of *A. chroococcum* melanin, as this feature has also been described for melanins synthesized by many different bacterial species (Bush, Simon, 2007). Microscopic analysis of *A. chroococcum* colonies revealed the presence of calcium crystals of various sizes in brown halo zones around the bacterial colonies, in the colonies themselves, and in the mucus. The crystals were absent or sparse in unpigmented colonies of this bacterial species.

Research conducted by Robson et al. (2015) showed the presence of 4 chromosomal genes (cum) encoding polyphenol oxidases in the genome of *A. chroococcum* strain NCIMB 8003. In addition, they found that the strain mentioned above of *A. chroococcum* is capable of producing carotenes in addition to melanin synthesis. Colonies of this strain on Burk's substrate took on a brown-black color, while on nutrient agar, they took on a yellow-brown color. The chromosome of *A. chroococcum* NCIMB 8003

contains 7 genes (crtZ, crtE, crtX, crtB, crtY, crtI, idi) encoding proteins, i.e., CrtE, CrtB, CrtY and CrtI involved in the synthesis of the carotene-like compound. The genes exhibit more than 60% identity with the orthologs of genes present in the genome of *Pseudomonas stutzeri*. At the same time, they were not detected in the genome of strain *A. vinelandii* DJ (Klassen, 2010). Melanins produced by *A. chroococcum* protect cells from hydroxyl radicals formed in the Fenton reaction, whereas carotenes are excellent antioxidants (Shivprasad, Page, 1989; Robson et al., 2015).

Literature data suggest that melanin, especially alloxanthin, has properties similar to those of soil humic substances, which promote the growth and yield of plants (Sutton, Sposito, 2005; Plonka, Grabacka, 2006; Muscolo et al., 2007). In their study, Gospodaryov, Lushchak (2011) observed the stimulating effect of purified melanin produced by *A. chroococcum* on such plants as lettuce, tomato, rapeseed and radish. The promotion of plant growth and development by melanin is also related to its ability to maintain water in the soil (Russo, Berlyn, 1990). In addition, the *A. chroococcum* strain is an excellent indicator of the presence of benzoic acid, and its salts in various substrates since precipitation of synthesized melanin occurs only in the presence of benzoic acid. Melanins extracted from *A. chroococcum* cultures, thanks to their affinity for metals and high adsorption capacity, are used in the bioremediation of soils and waters contaminated with heavy metals (Rizvi et al., 2019).

The research results available in the literature can contribute to clarifying the ecological role of melanins produced by soil bacteria of the genus *Azotobacter* and the importance of melanization processes in nature. Possible application of bacterial melanins in various industries

Numerous bacterial species, including pathogens, are capable of producing melanin. This pigment exerts a variety of functions, always beneficial to the host. Natural sources of melanin are currently being sought, among which melanins synthesized by microorganisms (bacteria, fungi) are the most promising. Their advantage is that they can be manufactured on a large scale, relatively cheaply, compared to the cost of obtaining synthetic melanins (Goncalvez, Pombeiro-Sponchiado, 2005). Melanins have been intensively studied for their properties and wide range of practical applications. Melanins produced by bacteria have been shown to effectively protect fibroblasts from UV radiation, indicating that they could be used as an ingredient in sunscreens (Geng et al., 2008; Kurian, Bhat, 2018). They have also found applications in medical diagnostics as a contrast agent in the optoacoustic tomography (Liopo et al., 2015). The pigments in question exhibit semiconductor properties and conduct electricity, hence the potential for their use in many branches of technology. Synthetic DOPA-melanin has been used for in situ formation of semiconductor coatings (Lee et al., 2007). There are also

sunglasses on the market with lenses containing melanin, which absorb ultraviolet radiation, thereby protecting vision (Lopusiewicz, Lisiecki, 2016). Melanin produced by *Shewanella oneidensis* increases electricity production in microbial fuel cells thanks to its ability to transport electrons (Turick et al., 2010). Research in recent years has drawn attention to the possibility of using melanins in innovative nanotechnologies, such as synthesising crystalline, flexible and thermostable nanocomposite films used in biomedicine (Kiran et al., 2017). The applicability of melanins in molecular biology is only feasible if pigment synthesis is regulated by a single gene or operon that serves as a marker for plasmid transfer from one organism to another. This is an excellent alternative to markers such as  $\beta$ -galactosidase or proteins responsible for antibiotic resistance (Tseng et al., 1990). From a medical point of view, melanogenic microorganisms are model microorganisms used to study the effects of various factors on melanin production. The anti-cancer properties of allomelanins are known (Kamei et al., 1997), as well as the inhibitory effect of melanins on HIV proliferation (Montefiori, Zhou, 1991; Sidibe et al., 1996). Melanin extracted from *Pseudomonas balearica* demonstrates antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans* (Zerrad et al., 2014). Thanks to their affinity for metals and high adsorption capacity, melanogenic bacteria can be used in bioremediation. It is recommended to detoxify soils or waters contaminated with heavy metals using melanin-producing microorganisms such as *Azotobacter chroococcum* (Rizvi et al., 2019) and *Pseudomonas stutzeri* (Thaira et al., 2019; Manirethan et al., 2018). In turn, melanin extracted from *E. coli* cultures has been used successfully for the bioremediation of drug-contaminated wastewater (Gustavsson et al., 2016).

## CONCLUSIONS

Melanins are compounds with unique attributes. The ability to produce the pigment group mentioned above is widespread among bacteria, and their role is complex and multifaceted. Melanins protect bacterial cells from external environmental factors, i.e. extreme temperatures, UV radiation, oxidizing agents, heavy metals and antibiotics. Melanins of natural origin synthesized by melanogenic microorganisms (bacteria, fungi) are increasingly being studied and applied. The advantage of natural melanins over synthetic ones is that they can be manufactured on a large scale by relatively simple and inexpensive methods. Melanins are intensively studied for their properties and wide range of practical applications. For example, obtaining high-performance microbial melanins involves selecting suitable microorganisms that can produce melanin extracellularly from an exogenous substrate and improving the metabolic process by adding tyrosine and copper to the culture substrate. Melanins produced by bacteria have found

applications in modern medicine, biotechnology, cosmetology and technology. From an agricultural perspective, these pigments can be used to promote plant growth and development and in the bioremediation of soils and waters contaminated with heavy metals. However, these pigments still hold many secrets and given their unique properties and role, they fully deserve continued research into new possibilities for their application.

## REFERENCES

- Agodi A., Stefani S., Corsaro C., Campanile F., Gribaldo S., Sichel G., 1996.** Study of a melanic pigment of *Proteus mirabilis*. *Research in Microbiology*, 147(3): 167-174.
- Ahmad S., Lee S.Y., Kong H.G., Jo E.J., Choi H.K., Khan R., Lee S.W., 2016.** Genetic determinants for pyomelanin production and its protective effect against oxidative stress in *Ralstonia solanacearum*. *PLOS ONE*, 11: e0160845, doi: 10.1371/journal.pone.0160845.
- Aquilanti L., Favilli F., Clementi F., 2004a.** Comparison of different strategies for isolation and preliminary identification of *Azotobacter* from soil samples. *Soil Biology and Biochemistry*, 36: 1475-1483.
- Aquilanti L., Mannazzu I., Papa R., Cavalca L., Clementi F., 2004b.** Amplified ribosomal DNA restriction analysis for the characterization of *Azotobacteraceae*: a contribution to the study of these free-living nitrogen-fixing bacteria. *Journal of Microbiological Methods*, 57: 197-206, doi: 10.1016/j.mimet.2004.01.006.
- Banerjee A., Supakar S., Banerjee R., 2014.** Melanin from the nitrogen-fixing bacterium *Azotobacter chroococcum*: A spectroscopic characterization. *PLOS ONE*, 9(1): e84574, doi: 10.1371/journal.pone.0084574.
- Baine W., Rasheed J.K., Feeley J.C., Gorman G.W., Casida L.E., 1978.** Effect of supplemental L-tyrosine on pigment production in cultures of the Legionnaire's disease bacterium. *Current Microbiology*, 1: 93-94.
- Beijerinck M.W., 1901.** Über ologonitrophile Mikroben. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. II Abt.*, 9: 561-582.
- Bortels H., Henkel H.G., 1968.** Über die Farbstoffbildung durch *Azotobacter chroococcum*. II. Das Chromogen. *Archiv für Mikrobiologie*, 60: 99-110.
- Bothma J.P., De Boor J., Divakar U., Schwenn P.E., Meredith P., 2008.** Device-quality electrically conducting melanin thin films. *Advanced Materials*, 20: 3539-3542, doi: 10.1002/adma.20070 3141.
- Bush W.D., Simon J.D., 2007.** Quantification of Ca<sup>2+</sup> binding to melanin supports the hypothesis that melanosomes serve a functional role in regulating calcium homeostasis. *Pigment Cell Research*, 20: 134-139.
- Chatfield C.H., Cianciotto N.P., 2007.** The secreted pyomelanin pigment of *Legionella pneumophila* confers ferric reductase activity. *Infection and Immunity*, 75: 4062-4070.
- Cheng J., Moss S.C., Eisner M., Zschack P., 1994.** X-ray characterization of melanins-I. *Pigment Cell Research*, 7: 255-262.
- Cordero R.J.B., Casadevall A., 2017.** Functions of fungal melanin beyond virulence. *Fungal Biology Reviews*, 31: 99-112, doi: 10.1016/j.fbr.2016.12.003.

- Coyne V.E., Al-Harathi L., 1992. Induction of melanin biosynthesis in *Vibrio cholera*. Applied and Environmental Microbiology, 58(9): 2861-2865.
- Crippa P.R., Horak V., Prota G., Svoronos P., Wolfram L., 1989. Chemistry of Melanins. pp. 234-323. The Alkaloids, vol. 36, Academic Press, New York.
- d'Ischia M., Pezzella A., Napolitano A. et al., 2015. Melanins and melanogenesis: from pigment cells to human health and technological applications. Pigment Cell and Melanoma Research, 28: 520-544, doi: 10.1111/pcmr.12393.
- De Smedth J., Bauwens M., Tytgat R., De Ley J., 1980. Intra- and intergeneric similarities of ribosomal ribonucleic acid cistrons of free-living, nitrogen-fixing bacteria. International Journal of Systematic Bacteriology, 30: 106-122.
- Döbereiner J., 1966. *Azotobacter paspali* sp. nov., umabactéria fixadora de nitrogênio na rizosfera dePas-palum. Pesquisa Agropecuária Brasileira, 1: 357-365.
- Döbereiner J., 1995. Isolation and identification of aerobic nitrogen-fixing bacteria from soil and plants. pp. 134-141. In: Methods in Applied Soil Microbiology and Biochemistry; Alef K., Nannipieri P., Academic Press, London.
- Drewnowska J.M., Zambrzycka M., Kalska-Szostko B., Fiedoruk K., Swiecicka I., 2015. Melanin-like pigment synthesis by soil *Bacillus weihenstephanensis* isolates from northeastern Poland. PLOS ONE, 10: e0125428, doi: 10.1371/journal.pone.0125428.
- Fedorow H., Tribl F., Halliday G., Gerlach M., Riederer P., Double K.L., 2005. Neuromelanin in human dopamine neurons: comparison with peripheral melanins and relevance to Parkinson's disease. Progress in Neurobiology, 75(2): 109-124.
- Funa N., Ohnishi Y., Fuji I., Shibuya M., Ebizuka Y., Horinouchi S., 1999. A new pathway for polyketide synthesis in microorganisms. Nature, 400: 897-899.
- Geng J., Tang W., Wan X., Zhou Q., Wang X-J., Shen P., Lei T-C., Chen X-D., 2008. Photoprotection of bacterial derived melanin against ultraviolet A-induced cell death and its potential application as an active screen. Journal of the European Academy of Dermatology and Venereology, 22(7): 852-857.
- Goncalvez R., Pombeiro-Sponchiado S. R., 2005. Antioxidant activity of the melanin pigment extracted from *Aspergillus nidulans*. Biological and Pharmaceutical Bulletin, 28: 1129-1131.
- Gospodaryov D.V., Lushchak V., 2011. Some properties of melanin produced by *Azotobacter chroococcum* and its possible application in biotechnology. Biotechnology, 4(2): 61-69.
- Gustavsson M., Hörnström D., Lundh S., Belotserkovsky J., Larsson G., 2016. Biocatalysis on the surface of *Escherichia coli*: melanin pigmentation of the cell exterior. Scientific Reports, 6: 36117, doi: 10.1038/srep36117.
- Harir M., Bellahcene M., Baratto M.C., Pollini S., Rossolini G.M., Trabalzini L., Fatarella E., Pogni R., 2018. Isolation and characterization of a novel tyrosinase produced by Sahara soil actinobacteria and immobilization on nylon nanofiber membranes. Journal of Biotechnology, 265: 54-64, doi: 10.1016/j.jbiotec.2017.11.004.
- Hawkins F.K., Johnston A.W., 1988. Transcription of a *Rhizobium leguminosarum* biovar *phaseoli* gene needed for melanin synthesis is activated by nifA of *Rhizobium* and *Klebsiella pneumoniae*. Molecular Microbiology, 2: 331-337.
- Herter S., Schmidt M., Thompson M.L., Mikolasch A., Schauer F., 2011. A new phenol oxidase produced during melanogenesis and encystment stage in the nitrogen-fixing soil bacterium *Azotobacter chroococcum*. Applied Microbiology and Biotechnology, 90: 1037-1049, doi: 10.1007/s00253-011-3093-x.
- Howey R.T., Lock C.M., Moore L.V.H., 1990. Subspecies names automatically created by Rule 46. International Journal of Systematic Bacteriology, 40: 317-319.
- Hynes M.F., Brucksch K., Priefer U., 1988. Melanin production encoded by a cryptic plasmid in a *Rhizobium leguminosarum* strain. Archives of Microbiology, 150: 326-332.
- Jacobson E.S., 2000. Pathogenic roles for fungal melanins. Clinical Microbiology Reviews, 13: 708-717.
- Jin H., Wang H., Zhang Y., Hu T., Lin Z., Liu B., Ma J., Wang X., Liu Q., Lin X., Xie Z., 2020. Description of *Azotobacter chroococcum* subsp. *isscasi* subsp. nov. isolated from paddy soil and establishment of *Azotobacter chroococcum* subsp. *chroococcum* subsp. nov. International Journal of Systematic and Evolutionary Microbiology, 70: 2124-2131, doi: 10.1099/ijsem.0.004026.
- Ju K.Y., Lee Y., Lee S., Park S.B., Lee J.K., 2011. Bioinspired polymerization of dopamine to generate melanin-like nanoparticles having an excellent free-radical-scavenging property. Biomacromolecules, 12: 625-632, doi: 10.1021/bm101281b.
- Kamei H., Koide T., Kojima T., Hasegawa M., Umeda T., 1997. Suppression of growth of cultured malignant cells by allomelanins, plant-produced melanins. Cancer Biotherapy and Radiopharmaceuticals, 12(1): 47-49.
- Karlsson O., Lindquist N.G., 2016. Melanin and neuromelanin binding of drugs and chemicals: toxicological implications. Archives of Toxicology, 90 :1883-1891, doi: 10.1007/s00204-016-1757-0.
- Keith K.E., Killip L., He P., Moran G.E., Valvano M.A., 2007. *Burkholderia cenocepacia* C5424 produces a pigment with antioxidant properties using a homogentisate intermediate. Journal of Bacteriology, 189: 9057-9065.
- Kimura T., Fukuda W., Sanada T., Imanaka T., 2015. Characterization of water-soluble dark-brown pigment from Antarctic bacterium, *Lysobacter oligotrophicus*. Journal of Bioscience and Bioengineering, 120: 58-61, doi: 10.1016/j.jbiosc.2014.11.020.
- Kiran G.S., Jackson S.A., Priyadharsini S., Dobson A.D.W., Selvin J., 2017. Synthesis of Nm-PHB (nanomelanin-polyhydroxy butyrate) nanocomposite film and its protective effect against biofilm-forming multi drug resistant *Staphylococcus aureus*. Scientific Reports, 7: 9167, doi: 10.1038/s41598-017-08816-y.
- Klassen J.L., 2010. Phylogenetic and evolutionary patterns in microbial carotenoid biosynthesis are revealed by comparative genomics. PLOS ONE, 5(6): e11257, doi: 10.1371/journal.pone.0011257.
- Kotob S., Coon S.I., Quintero E.J., Weiner R.M., 1995. Homogentisic acid is the primary precursor of melanin synthesis in *Vibrio cholerae*, a *Hyphomonas* strain and *Shewanella colwelliana*. Applied and Environmental Microbiology, 61: 1620-1621.
- Krasil'nikov N.A., 1949. Guide to the bacteria and actinomycetes. Red. Akademia Nauk SSSR, Moscow.
- Kurian N.K., Bhat S.G., 2018. Data on the characterization of noncytotoxic pyomelanin produced by marine *Pseudomonas stutzeri* BTCZ10 with cosmetological importance. Data in Brief, 18: 1889-1894, doi: 10.1016/j.dib.2018.04.123.

- Langfelder K., Streibel M., Jahn B., Haase G., Brakhage A.A., 2003.** Biosynthesis of fungal melanins and their importance for human pathogenic fungi. *Fungal Genetics and Biology*, 38: 143-158.
- Lee H., Dellatore S.M., Miller W.M., Messersmith P.B., 2007.** Mussel-inspired surface chemistry for multifunctional coatings. *Science*, 318: 426-430.
- Le Na N.T., Duc Loc S., Minh Tri N.L. et al., 2019.** Nanomelanin potentially protects the spleen from radiotherapy-associated damage and enhances immunoactivity in tumor-bearing mice. *Materials*, 12: 1725, doi: 10.3390/ma12101725.
- Lenart A., 2012.** Occurrence, characteristics, and genetic diversity of *Azotobacter chroococcum* in various soils of southern Poland. *Polish Journal of Environmental Studies*, 21(2): 415-424.
- Lin W.P., Lai H.L., Liu Y.L., Chiung Y.M., Shiao Ch.Y., Han J.M., Yang Ch.M., Liu Y.T., 2005.** Effect of melanin produced by recombinant *Escherichia coli* on antibacterial activity of antibiotics. *Journal of Microbiology, Immunology and Infection*, 38: 320-326.
- Liopo A., Su R., Oraevsky A., 2015.** Melanin nanoparticles as a novel contrast agent for optoacoustic tomography. *Photoacoustics*, 3: 35-43.
- Lipman J.G., 1903.** Experiments on the transformation and fixation of nitrogen by bacteria. Report on the New Jersey Agricultural Experiment Station, 24: 217-285.
- Lipman J.G., 1904.** Soil bacteriological studies. Further contributions to the physiology and morphology of members of the *Azotobacter* group. Report on the New Jersey Agricultural Experiment Station, 25: 237-289.
- Liu G.Y., Nizet V., 2009.** Color me bad: microbial pigments as virulence factors. *Trends in Microbiology*, 17:406-413.
- Liu L., Yuan T., An Q., Yang M., Mao X., Mo C., Tan Z., Peng G., 2019.** *Azotobacter bryophylli* sp. nov., isolated from the succulent plant *Bryophyllum pinnatum*. *International Journal of Systematic and Evolutionary Microbiology*, 69: 1986-1992. doi: 10.1099/ijsem.0.003412
- Lopez-Serrano D., Solano F., Sanchez-Amat A., 2004.** Identification of an operon involved in tyrosinase activity and melanin synthesis in *Marinomonas mediterranea*. *Gene*, 342: 179-187.
- Lopusiewicz L., Lisiecki S., 2016.** „Black gold” – melanins in human life. *Kosmos*, 65(4): 621-629. (in Polish + summary in English)
- Manirethan V., Raval K., Rajan R., Thaira H., Balakrishnan R.M., 2018.** Data on the removal of heavy metals from aqueous solution by adsorption using melanin nanopigment obtained from marine source: *Pseudomonas stutzeri*. Data in Brief, 20: 178-189, doi: 10.1016/j.dib.2018.07.065.
- Martyniuk S., Martyniuk M., 2003.** Occurrence of *Azotobacter* spp. in some Polish soils. *Polish Journal of Environmental Studies*, 12: 371-374.
- Montefiori D.C., Zhou J.Y., 1991.** Selective antiviral activity of synthetic soluble L-tyrosine and L-dopa melanins against human immunodeficiency virus in vitro. *Antiviral Research*, 15(1): 11-15.
- Musco A., Sidari M., Francioso O., Tugnoli V., Nardi S., 2007.** The auxin-like activity of humic substances is related to membrane interactions in carrot cell cultures. *Journal of Chemical Ecology*, 33(1): 115-129.
- Nappi A., Ottaviani E., 2000.** Cytotoxicity and cytotoxic molecules in invertebrates. *BioEssays*, 22: 469-480.
- Nicolaus R.A., Piattelli M., Fattorusso E., 1964.** The structure of melanins and melanogenesis. IV: On some natural melanins. *Tetrahedron*, 20: 1163-1172.
- Nicolaus R., 1968.** Melanins. Paris, Hermann.
- Noorian P., Hu J., Chen Z., Kjelleberg S., Wilkins M.R., Sun S., McDougald D., 2017.** Pyomelanin produced by *Vibrio cholerae* confers resistance to predation by *Acanthamoeba castellanii*. *FEMS Microbiology Ecology*, 93(12), doi: 10.1093/femsec/fix147.
- Nosanchuk J.D., Casadevall A., 2003.** The contribution of melanin to microbial pathogenesis. *Cellular Microbiology*, 5: 203-223, doi: 10.1046/j.1462-5814.2003.00268.x.
- Nosanchuk J.D., Casadevall A., 2006.** Impact of melanin on microbial virulence and clinical resistance to antimicrobial compounds. *Antimicrobial Agents and Chemotherapy*, 50: 3519-3528, doi: 10.1128/AAC.00545-06.
- Nosanchuk J.D., Stark R.E., Casadevall A., 2015.** Fungal melanin: what do we know about structure? *Frontiers of Microbiology*, 6: 1463, doi: 10.3389/fmicb.2015.01463.
- Özen A.I., Ussery D.W., 2012.** Defining the *Pseudomonas* genus: where do we draw the line with *Azotobacter*? *Microbial Ecology*, 63: 239-248, doi: 10.1007/s00248-011-9914-8.
- Page W., Shivprasad S., 1991.** *Azotobacter salinestris* sp. nov., a sodium-dependent, microaerophilic, and aeroadaptive nitrogen-fixing bacterium. *International Journal of Systematic Bacteriology*, 41: 369-376.
- Paget M. S., Hintermann G., Smith C. P., 1994.** Construction and application of streptomycete promoter probe vectors which employ the *Streptomyces glaucescens* tyrosinase-encoding gene as a reporter. *Gene*, 146: 105-110.
- Paul E.A., Clark F.E., 2000.** *Mikrobiologia i biochemia gleb.* Jędrych M., Wydawnictwo Uniwersytetu Marii Curie-Skłodowskiej, Lublin, 400 pp.
- Pavan M.E., Abbott S.L., Zorzópulos J., Janda J.M., 2000.** *Aeromonas salmonicida* subsp. *pectinolytica* subsp. nov., a new pectinase-positive subspecies isolated from a heavily polluted river. *International Journal of Systematic and Evolutionary Microbiology*, 50: 1119-1124, doi: 10.1099/00207713-50-3-1119.
- Patel K., Wyman J., Patel K., Burden B., 1996.** A mutant of *Bacillus thuringiensis* producing a dark-brown pigment with increased UV resistance and insecticidal activity. *Journal of Invertebrate Pathology*, 67: 120-124.
- Piattelli M., Fattorusso E., Nicolaus R.A., Magno, S., 1965.** The structure of melanins and melanogenesis. V. Ustilago melanin. *Tetrahedron*, 21: 3229-3236.
- Piñero S., Rivera J., Romero D., Cevallos M.A., Martínez A., Bolívar F., Gosset G., 2007.** Tyrosinase from *Rhizobium etli* is involved in nodulation efficiency and symbiosis-associated stress resistance. *Journal of Molecular Microbiology and Biotechnology*, 13: 35-44, doi: 10.1159/000103595.
- Plonka P.M., Grabacka M., 2006.** Melanin synthesis in microorganisms- biotechnological and medical aspects. *Acta Biochimica Polonica*, 53(3): 429-443.
- Pralea I.E., Moldovan R.C., Petrache A.M. et al., 2019.** From extraction to advanced analytical methods: the challenges of melanin analysis. *International Journal of Molecular Sciences*, 20: 3943, doi: 10.3390/ijms20163943.
- Prabhakaran K., Kirchheimer W.F., Harris E.B., 1968.** Oxidation of phenolic compounds by *Mycobacterium leprae* and inhibition of phenolase by substrate analogs and copper chelators. *Journal of Bacteriology*, 95: 2051-2053.



- Prota G., 1992.** Melanins and melanogenesis. San-Diego, Academic Press.
- Przemysław M.P., Maja G., 2006.** Melanin synthesis in microorganisms: Biotechnological and medical aspects. *Acta Biochimica Polonica*, 53: 429-443.
- Rizvi A., Ahmed B., Zaidi A., Khan M.S., 2019.** Bioreduction of toxicity influenced by bioactive molecules secreted under metal stress by *Azotobacter chroococcum*. *Ecotoxicology*, 28: 302-322, doi: 10.1007/s10646-019-02023-3.
- Robson R.L., Jones R., Robson R.M., Schwartz A., Richardson T.H., 2015.** *Azotobacter* genomes: The genome of *Azotobacter chroococcum* NCIMB 8003 (ATCC 4412). *PLOS ONE*, 10: 1-35, doi: 10.1371/journal.pone.012799.
- Rubio E.J., Montecchia M.S., Tosi M., Cassán F.D., Peticari A., Correa O.S., 2013.** Genotypic characterization of *Azotobacteria* isolated from Argentinean soils and plant-growth-promoting traits of selected strains with prospects for biofertilizer production. *The Scientific World Journal*, pp. 1-12, doi: 10.1155/2013/519603.
- Russo R.O., Berlyn G.P., 1990.** The use of organic biostimulants to help low input sustainable Agriculture. *Journal of Sustainable Agricultural*, 1(2): 19-42.
- Sadasivan L., Neyra C.A., 1987.** Cyst production and brown pigment formation in aging cultures of *Azospirillum brasilense* ATCC 29145. *Journal of Bacteriology*, 169: 1670-1677.
- Saud H.M., Alaubiydi M.A., 2016.** Production, extraction and partial purification of melanin pigment from pathogenic *Klebsiella pneumoniae* HM isolated from clinical samples. *International Journal of Current Microbiology and Applied Sciences*, 5: 910-919.
- Shivprasad S., Page W.J., 1989.** Catechol formation and melanization by Na<sup>+</sup>-dependent *Azotobacter chroococcum*: a protective mechanism for aeroadaptation? *Applied and Environmental Microbiology*, 55(7): 1811-1817, doi: 10.1128/aem.55.7.1811-1817.1989.
- Sidibe S., Saal F., Rhodes-Feuillet A., Lagaye S., Pelicano L., Canivet M., Peries J., Dianoux L., 1996.** Effects of serotonin and melanin on in vitro HIV-1 infection. *Journal of Biological Regulators and Homeostatic Agents*, 10(1): 19-24.
- Solano F., 2014.** Melanins: skin pigments and much more. Types, structural models, biological functions, and formation routes. *New Journal of Science*, pp. 1-28, doi: 10.1155/2014/498276.
- Stolarzewicz I., Kapturowska A., Bialecka-Florjańczyk E., 2012.** Mikrobiologiczne źródła barwników w technologii żywności. *Postępy Mikrobiologii*, 51: 167-176.
- Sutton R., Sposito G., 2005.** Molecular structure of soil humic substances: the new view. *Environmental Science and Technology*, 39(3): 9009-9015.
- Tchan Y.T., New P.B., 1984.** *Azotobacteraceae*. pp. 220-229. In: *Bergey's Manual of Systematic Bacteriology* vol. 1; Kreig, N.R., Holt J.G., Williams and Wilkins, Baltimore.
- Thaira H., Raval K., Manirethan V., Balakrishnan R.M., 2019.** Melanin nano-pigments for heavy metal remediation from water. *Separation Science and Technology*, 54: 265-274, doi: 10.1080/01496395.2018.1443132.
- Thompson J.P., Skerman V.B.D., 1979.** *Azotobacteraceae*. p. 277. In: *The Taxonomy and Ecology of the Aerobic Nitrogen-Fixing Bacteria*. Academic Press London, New York, Toronto, Sydney, San Francisco.
- Tran-Ly A.N., Ribera J., Schwarze F.W.M.R., Brunelli M., Fortunato G., 2020.** Fungal melanin-based electrospun membranes for heavy metal detoxification of water. *Sustainable Materials and Technologies*, 23: e00146, doi: 10.1016/j.susma.t.2019.e00146.
- Tran-Ly A.N., Reyes C., Schwarze F.W.M.R., Ribera J., 2020.** Microbial production of melanin and its various applications. *World Journal of Microbiology and Biotechnology*, 36: 1-9, <https://doi.org/10.1007/s11274-020-02941-z>.
- Tseng H.C., Lin C.K., Hsu B.J., Leu W.M., Lee Y.H., Chiou S.J., Hu N.T., Chen C.W., 1990.** The melanin operon of *Streptomyces antibioticus*: expression and use as a marker in gram-negative bacteria. *Gene*, 86: 123-128.
- Turick C.E., Caccavo F.Jr., Tisa L.S., 2008.** Pyomelanin is produced by *Shewanella algae* BrY and affected by exogenous iron. *Canadian Journal of Microbiology*, 54(4): 334-339.
- Turick C.E., Knox A.S., Becnel J.M., Ekechukwu A.A., Milliken C.E., 2010.** Properties and function of pyomelanin. 449-472. Elnashar M., *Biopolymers*, 1st edn. Sciyo, Rijeka, doi: 10.5772/10273.
- Valeru S.P., Rompikuntal P.K., Ishokawa T., Vaitkevicius K., Sjoling A., Dolganov N., Zhu J., Schoolink G., Wai S.N., 2009.** Role of melanin pigment in expression of *Vibrio cholerae* virulence factors. *Infection and Immunity*, 77: 935-942.
- Wolska K.I., Grudniak A.M., Kraczkiewicz-Dowjat A., Kurek A., 2010.** Różnorodne funkcje wybranych pigmentów bakteryjnych. *Postępy Mikrobiologii*, 40: 105-114.
- Zeng Z., Guo X.P., Cai X., Wang P., Li B., Yang J.L., Wang X., 2017.** Pyomelanin from *Pseudoalteromonas lipolytica* reduces biofouling. *Microbial Biotechnology*, 10: 1718-1731, doi: 10.1111/1751-7915.12773.
- Zerrad A., Anissi J., Ghanam J., Sendide K., El Hassouni M., 2014.** Antioxidant and antimicrobial activities of melanin produced by a *Pseudomonas balearica* strain. *Biotechnology Letters*, 5: 87-94.
- Zughaier S.M., Ryley H.C., Jackson S.K., 1999.** A melanin pigment purified from an epidemic strain of *Burkholderia cepacia* attenuates monocyte respiratory burst activity by scavenging superoxide anion. *Infection and Immunity*, 67: 908-913.

Author

ORCID

Monika Kozieł

0000-0001-7653-3610

received – 5 October 2022

revised – 11 December 2022

accepted – 30 December 2022

The author declare no conflict of interest.



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike (CC BY-SA) license (<http://creativecommons.org/licenses/by/4.0/>).