

An attempt to assess the usefulness of Google Scholar as a data source for entomological review papers on the example of the description of the biology of *Anomala dubia* (Scopoli, 1763) (Coleoptera: Scarabaeidae)

Alina Bochniarz

Department of Extension and Information
 Institute of Soil Science and Plant Cultivation – State Research Institute
 ul. Czarzoryskich 8, 24-100 Puławy, POLAND

Abstract. Review papers, especially those with features of meta-analysis, are a valuable part of the scientific literature, both from the point of view of specialists and researchers seeking information in fields new to them. The basis of this type of publication is the proper selection of study sources. The aim of this study was to assess the usefulness of the Google Scholar search engine in the selection of material for review papers in entomology. The analysis was based on a sample assignment involving the characterisation of the beetle species *Anomala dubia*. Problems arising in using the retrieved resource were identified and assigned to selected categories. The search results included almost all relevant records from the SCOPUS and Web of Science databases, plus a number of potentially valuable publications from other sources. The collected material was not sufficient to characterise the studied species in detail, but this could also be due to the lack of digitalised data in the available literature resources.

Keywords: Google Scholar, *Anomala dubia*, scientific publications, bibliographic data

INTRODUCTION

The widespread use of the Internet has changed the accessibility of scientific papers. Many services and repositories have been created where they are collected (e.g. the Polish Repository of Centre for Open Science, repositories of universities, Zenodo), it is also connected with green way of open access. Google Scholar is a free scientific search engine that is widely available, also to scientists from developing countries, covering about half of the world's scientific articles in open access (Singh et al., 2023). For biology and chemistry, it is ranked first in the world (Klopfenstein, Dampier, 2021). It is estimated to include 389 million records (Gusenbauer, Haddaway, 2020). The filters offered, although considered insufficient (What

are the advantages ...), allow a certain degree of source ordering, which facilitates initial orientation in the extensive resources. In addition, Google's online translators make it possible to understand texts in numerous rare national languages. The work on the use of Google Scholar in science is mainly concerned with the usefulness of the biblio- and scientometric indicators offered by the search engine (e.g. Pereira, Mugnaini, 2023) and its comparison with database resources and the capabilities of other search engines (e.g. Ungar et al., 2023). Prestigious bibliographic databases such as SCOPUS or Web of Science only include journal articles that meet strict criteria, which, at least in theory, guarantees their high scientific quality. Google Scholar has a significant quantitative advantage over these databases because, in addition to peer-reviewed scientific papers with an international presence, it includes local scientific publications, conference proceedings, research reports, textbooks, book chapters, patents, etc.

However, the diversity of origins of Google Scholar searches can be a source of problems that limit the usefulness of the material so retrieved (What are the advantages...). The aim of this paper is to try to identify such difficulties and their causes. The analysis was carried out using the example of information on the beetle *Anomala dubia*. It is probably the only representative of the genus in Poland (Burakowski et al., 1983), often, even in scientific publications, confused with other representatives of its family (Thomaes et al., 2022; Urban et al., 2020). It is not one of the economically important pests of agricultural crops, but information about it can be useful in agroforestry or energy crop production. It deserves attention due to its affinity with species of greater economic importance that may soon reach Poland (PIORIN, 2016).

MATERIAL AND METHOD

The material for analysis consisted of all records retrieved in March 2023 from Google Scholar in response to the query 'Anomala dubia', with no time limit. The pub-

Corresponding author:

Alina Bochniarz
 e-mail: aboch@iung.pulawy.pl
 phone +48 81 4786 726




Figure 1. The study scheme.

lications, if it was possible, were searched for the word ‘Anomala’, as well as ‘dubia’, to also include the use of the abbreviated species name (*A. dubia*). The paragraphs contained key words, if it was necessary to understand, were translated independently (Russian, Ukrainian, English, French) or using appropriate Google translators. Problems that arose when using the resource were identified and categorised into one of the distinguished groups: technical, bibliographical or substantive (Fig. 1). Technical problems included difficulties in opening or searching the files (files that cannot be opened, uncopyable or unsearchable files, additional copies). Records under technical problems were excluded from the further study. Bibliographical problems related mainly to the structure of the works and shortcomings in the bibliographic data. Substantive problems referred to the quality of the presented data. In addition, a list of the most important issues in the resource acquired in the manner described is presented. The review of issues and the literature list omit items relating only to the recording of the presence of *A. dubia* in the study area (“simple checklists” in Fig. 1). Similar search (“Anomala dubia”, all open access, no time limits, institutional access) was conducted in SCOPUS and Web of Science databases.

RESULTS AND DISCUSSION

Types of publication

The selected collection of records consisted mainly of lists of species found in different areas. These covered a diverse range of organisms: e.g. protected insects, beetles in general or only from the Scarabaeidae family. The search results also included papers published in scientific journals. It should be noted that almost all open-access published articles on the selected topic collected in the prestigious SCOPUS and Web of Science databases were included, but the only item omitted was an important study on the morphology and biology of larvae¹. Depending on

¹ Micó E., Galante E., 2005. Larval morphology and biology of some European Anomalini (Coleoptera: Scarabaeoidea: Rutelidae: Anomalinae). A phylogenetical approach. *Insect Systematics and Evolution*, 36: 183-198. It appeared in the results of the search performed in the same way in April 2023.

the topic studied and the criteria adopted, the percentage of coverage of results from Google Scholar and the databases compared to it varies, but can reach 100 (Ungar et al., 2023). In addition to this, the resource included master’s and doctoral theses, research project reports, conference proceedings, reports from scientific trips and meetings, posters, and book chapters. Such a wide range of included sources is characteristic of Google Scholar (Racki, Drabek, 2013) and is assessed as both a disadvantage and an advantage. With regard to *A. dubia*, it was a source of much interesting information that did not appear in studies dealing directly with the characterisation of this species, for example related to the presence of *A. dubia* in the diet of different bird species. Access to additional literature, including little-known contributory works, can sometimes be inspiring and allows new research directions to be identified. However, at the same time, it forces a critical approach to the data used, as not all sources have been verified through the scientific review process.

Overview of key issues in selected material

The chapter gives the most important issues that are usually discussed when characterising an insect species, as well as additional information. In many publications, reference to *A. dubia* was merely in the nature of citing data from other sources. The records covered the period of 1936–2022.

• TAXONOMY

Audras et al., 1953; Cosandey et al., 2017; Dikmen, Özuluğ, 2018; Greven, 2011; Hielkema, 2022; Kreil, 1993; Kuntze, 1936

• EXTERNAL MORPHOLOGY

Description and photographs: Charrier, 2002; Frantsevich, 2004; Horvath et al., 2014; Petrova et al., 2022

Origin of the Latin name: Forshage, 2000

Body size, weight: Orłowski et al., 2011; Pedersen et al., 2012; Watkins, Mann, 2018a; Wouters, van Oosten, 2013

Table 1. *Anomala dubia* host-plants acc. to literature.

Host-plant (Latin name)	Imago	Source	Host-plant (Latin name)	Source
<i>Acer campestre</i>	Horvath et al., 2014		<i>Rosa canina</i>	Horvath et al., 2014
<i>Acer negundo</i>	Dumas, 2019		<i>Rubus</i>	Kuntze, 1936
<i>Alnus glutinosa</i>	Petrova et al., 2022		<i>Salix</i>	Audras et al., 1953; Delwaide, Thieren, 2010; Gallou, 1985; Georgi, Müller, 2015; Kühne et al., 2000; Marczak, Mroczynski, 2016; Mertlik, 2011; Nicolas, Nicolas, 1965; Olsson, Lemdahl, 2009; Teodorescu, Maican, 2014
<i>Betula</i>	Gallou, 1985; Olsson, Lemdahl, 2009		<i>Salix daphnoides</i>	Berglind et al., 1999
<i>Betula pendula</i>	Horvath et al., 2014		<i>Salix nigricans</i>	Berglind et al., 1999
<i>Castanea sativa</i>	Chittaro, Sanchez, 2015; Horvath et al., 2014		<i>Salix triandra</i>	Berglind et al., 1999
<i>Cirsium</i>	Teodorescu, Maican, 2014		<i>Sophora japonica</i>	Horvath et al., 2014
<i>Corylus</i>	Kuntze, 1936		<i>Sorbus aria</i>	Horvath et al., 2014
<i>Crataegus monogyna</i>	Horvath et al., 2014		<i>Sorbus aucuparia</i>	Horvath et al., 2014
<i>Euphorbia</i>	Teodorescu, Maican, 2014		<i>Tilia cordata</i>	Horvath et al., 2014
<i>Fagus</i>	Bitokon' et al., 2005		<i>Ulmus campestris</i>	Horvath et al., 2014
<i>Hippophaë rhamnoides</i>	Reiter, 1986		<i>Taccinium corymbosum</i>	Janiuk, Gantner, 2012
<i>Malus</i>	Kuntze, 1936		<i>Vitis</i>	Audras et al., 1953; Teodorescu, Maican, 2014; Voigt, Tóth, 2004
<i>Pinus</i>	Gutowski et al., 2006; Kavka et al., 2022; Kuntze, 1936; Prakhodskiy et al., 2010; Veremeev, 2005			
<i>Platanus acerifolia</i>	Horvath et al., 2014			
<i>Populus</i>	Georgi, Müller, 2015; Horvath et al., 2014; Radzikowski et al., 2020; Teodorescu, Maican, 2014		Larva	
<i>Populus alba</i>	Meleshko, 2003		<i>Acacia</i>	Netou et al., 2007
<i>Populus nigra</i>	Horvath et al., 2014		<i>Ammophila arenaria</i> (grass)	de Rond, 2016
<i>Populus simonii</i>	Meleshko, 2003		<i>Beta vulgaris</i>	Spaar et al., 1988
<i>Populus tremula</i>	Meleshko, 2003		<i>Fagopyrum</i>	Naumkin, Lysenko, 2014
<i>Prunus avium</i>	Horvath et al., 2014		<i>Festuca</i>	Mertlik, Brůha, 2011
<i>Prunus persica</i>	Voigt, Tóth, 2004, 2009		<i>Fraxinus</i>	Netou et al., 2007
<i>Prunus persica</i> var. <i>nucipersica</i>	Voigt, Tóth, 2004, 2009		<i>Gleditsia</i>	Netou et al., 2007
<i>Rhamnus</i>	Teodorescu, Maican, 2014		<i>Pinus</i>	Kozel, 2012
<i>Rhus typhina</i>	Horvath et al., 2014		<i>Salix</i>	Netou et al., 2007
			<i>Solanum tuberosum</i>	Spaar et al., 1987

Table 2. Birds species in which diet *Anomala dubia* was recorded acc. to literature.

Bird (Latin name)	Source
<i>Coracias garrulus</i>	Hebda et al., 2019
<i>Erithacus rubecula</i>	Chaplygina et al., 2016
<i>Ficedula albicollis</i>	Chaplygina et al., 2015, 2022; Leženina et al., 2011
<i>Lanius collurio</i>	Goławski, 2006; Maréchal, 1993; Morelli et al., 2015; Nijssen, Geertsma, 2014; Pedersen et al., 2012; Tryjanowski et al., 2003; van der Kruis, 2008; van Duinen et al., 2004, 2005
<i>Larus cachimans</i>	Atamas', Loparev, 2015
<i>Motacilla flava</i>	Prokof'eva, 2013
<i>Oenanthe isabellina</i>	de Rond, 2016
<i>Oenanthe oenanthe</i>	van Oosten et al., 2008, 2014
<i>Phoenicurus ochruros</i>	Orłowski et al., 2011
<i>Saxicola rubetra</i>	Prokof'eva, 2013
<i>Saxicola rubicola</i>	van Oosten, van den Burg, 2015
<i>Turdus philomelos</i>	Harčenko et al., 2002
<i>Turdus viscivorus</i>	Prokof'eva, 2003, 2007
<i>Tyto alba</i>	De Pablo, 2000
<i>Upupa epops</i>	Nuhlíčková et al., 2016

variation in body colour: Décobert, 2010; Ostrovskij, 2015; Petrova et al., 2022; Telfer, Eversham, 1995.

Problems of species determination: Schacht, 2022; Thomaes et al., 2016, 2022; Urban et al., 2020

Circular polarization: Horvath et al., 2014.

Morphology and body size of larvae, species determination of *A. dubia* larvae: Laznik, Trdan, 2013; Maksimova, 2008.

- ANATOMY

Bohac et al., 2020; Frantsevich et al., 2014; Pacheco et al., 2022; Romero-Lopez et al., 2011

- OCCURRENCE AND DISTRIBUTION

Zoogeographic regions, subspecies: Chenchouni et al., 2015; Japoshvili et al., 2022; Smith, 2009; Šohin, 2011

Altitude a.s.l.: Mertlik, 2011; Polat et al., 2018; Szafraniec et al., 2021

Habitats: Aleksandrowicz et al., 2004; Berglind et al., 1999; Borowski et al., 2016; Egorov et al., 2017; Gallou, 1985; Lammerts et al., 2009; Marczak, Mroczynski, 2016; Martikainen, 2016; Mertlik, 2011; Ødegaard et al., 2001; Pedersen et al., 2012; Sažnev, Anikin, 2021; Schacht, 2017; Sörensson, 2020; Telfer, Eversham, 1995; Toth et al., 1994; van Duinen et al., 2004, 2005; Veremeev, 2005; Veremeev et al., 2001; Watkins, Mann, 2018a, 2018b; Wolender, Zych, 2007; Zhukov, 2006, 2007; Zúber, 2011

Anthropogenic habitats: Borowski et al., 2016; Delwaide, Thieren, 2010; Galinovskij et al., 2019; Jäger et al., 2014; Komaromi, 2020; Kubicka, 1981; Kušková, 2014

Types of habitats: Kühne et al., 2000; Nadvorny, 2007; Olsson, 2006; Ostrovskij, 2015

Changes in size and range of population, threat factors:

Bayr, Svae Johansen, 2022; Carpaneto et al., 2011; Kavka et al., 2022; Köteles, Bakonyi, 1996; Kubicka, 1981; Leontiev, 2021; Malicky, 1975; Ødegaard et al., 2001; Zúber, 2011

- DEVELOPMENT

Life cycle, imago flying period, egg deposition, factors affecting length of development: de Rond, 2016; Laznik, Trdan, 2013; Mertlik, 2011; Negrobov, 2009; Romero-Lopez et al., 2011; Ruta et al., 2016; Spaar et al., 1987, 1988; van Duinen et al., 2005; Voigt, Tóth, 2009

- HABITS

Non-feeding habitats, role in plant pollination: Bellmann, 2002; Dutto, 2007; Egorov et al., 2017; Jakubská-Busse, Kadej, 2008, 2011; Jakubská, Kadej, 2006; Leontiev, 2021; Radzikowski et al., 2020; Sažnev, Anikin, 2021

Trapping of insects, effect of polarized light: Bayr, Svae Johansen, 2022; Bukejs, 2006; Chenchouni et al., 2015; Horvath et al., 2014; Kofler, 1999; Martikainen, 2016; Radzikowski et al., 2020; Toth et al., 1994, 1997.

Insect motion: Frantsevich, 2004; Mertlik, 2011; Watkins, Mann, 2018b; Zúber, 2011.

Depth of larvae feeding in soil, larval adaptation to unfavourable conditions: Netoiu et al., 2007; van Duinen et al., 2005; Von Niklas, 1964

- FEEDING HABITS, HOST-PLANTS, FEEDING DAMAGE

Feeding habits of imago: Gutowski et al., 2006; Janiuk, Gantner, 2012; Kuntze, 1936; Voigt, Tóth, 2004.

Host-plants: Table 1.

Outbreaks, harmfulness, damage description: Baban, 2006; Georgi, Müller, 2015; Jameson et al., 2003; Kolk, Grodzki, 2011; Kuntze, 1936; Voigt, Tóth, 2004

Feeding on grass (dunes, grasslands), feeding on field crops, feeding in forests: de Rond, 2016; Kulikov, Naumkin, 2004; Laznik, Trdan, 2015; Maleshko, 2003; Netoiu et al., 2007; Radics, Mikóházi, 2010; Spaar et al., 1987, 1988; Voigt, Tóth, 2004

- RELATIONSHIPS WITH OTHER ORGANISMS

Predatory insects, spiders, parasitic insects: Amolin, 2005; Gusenleitner et al., 2008; Heijerman, Turin, 2017; Kavka et al., 2022; Reitze, Nentwig, 1991; Scharfy, 2012; Teppner, 2008; Von Klausnitzer et al., 2013; Von Niklas, 1964

Birds: Table 2

mammals: Lanszka et al., 2009; Laznik, Trdan, 2014; Mikheyev, 2002

parasitic nematodes: Garcia del Pino, Palomo, 1995; Gradinarov, Petrova, 2011; Gradinarov et al., 2011; Grewal et al., 1996; Han et al., 2012; Jarry, 1964; Nguyen, Smart, 1993; Parra, 2020; Peters, Galarza, 2004; Poinar, Kozodoi, 1988; Quintero Marin, 2003; Uhan, 2005; Wu, 2013

bacteria, protozoa, fungi: Bajc, 2016; Fătu et al., 2016, 2017; Kolasa et al., 2018; Laznik, Trdan, 2015; Lila, 2014; Niklas, 1963; Pandey, 2013; Théodoridès, 1963

- GENETICS, PHYLOGENESIS

Hendrich et al., 2015; Jameson et al., 2007; Ramírez-Ponce, Morón, 2009

- OTHER TOPICS

Content of F, Cl, Br, and I in larval body: Pokarzhevskii, Zhulidov, 1995

PROBLEMS IDENTIFIED IN THE USE OF SEARCH RESULTS

Technical problems

Among the files retrieved, several, despite the pdf extension, could not be opened, were probably corrupted or required non-standard software to view. Records in which the text was presented as an image or is uncopyable also posed a problem, making direct searching and copying into translators impossible. In the case of short documents in known languages, it is possible to search for interesting passages on your own, but in extensive publications this is very time-consuming. There were also duplicate records, which are not protected against by the search algorithm used by Google Scholar (What are the advantages ...). Two publications did not meet the criterion of searching, because words *Anomala* and *dubia* occurred only separately

and texts concerned to other animals. In the overall collection, the problematic files accounted for a small proportion. In the search carried out, there were no difficulties in accessing the full texts of publications, although according to user feedback, such problems may appear in results derived from Google Scholar (What are the advantages ...).

Bibliographical problems

Bibliographical problems are not due to the substantive quality of the publication, but make it difficult to use and cite. Due to the multitude of non-unified sources and the automatic processing of data, Google Scholar is particularly prone to such errors (Racki, Drabek, 2013; Sauvayre, 2022). Therefore, problems with the quality and transparency of bibliographic data are often reported (Doğan, 2022).

Also in the results of the search carried out, many records lacked the full bibliographic details of the publication. For example, only the title and author were given, while the page range indicated that it was part of a larger whole. It is possible to find missing data in repositories and citations, but this is time-consuming. This problem occurs most often with Russian and Ukrainian publications.

Different spellings of the names of the same authors also lead to citation errors. It may be due to the use of different tables or different ways of transliterating Cyrillic characters, and a similar problem is posed by the prepositions *de*, *von*, *van* before names.

In search results, there were publications in different languages: Bosnian, Croatian, Czech, Danish, Dutch, English, Finnish, French, German, Hungarian, Indonesian, Italian, Korean, Lithuanian, Norwegian, Polish, Portuguese, Romanian, Russian, Slovak, Slovenian, Spanish, Swedish, Ukrainian. English is generally known for scientists, but other languages can be a barrier for texts to be used by foreigners. In such cases, the lack of an English title and abstract makes it impossible to quickly select material for relevance. It is a good practice, relatively uncommon however, to also include English captions for tables and figures.

The lack of a single explanation of the abbreviations and symbols used in the text is also a handicap. Even if they are obvious to the author, readers using other languages may not properly understand them.

The use of, often very arbitrary, common names for animals and plants, different even in articles in the same language, leads to confusion. In the case of living organisms, the Latin nomenclature should be the standard in scientific papers, preferably used throughout the text, not just at the first occurrence of a name.

In the results of the search carried out, several examples can be found of publishing virtually the same content under different titles or with a different author composition. This is sometimes justified by a change of version depending on the target audience or is the result of artificial

over-publication by the author. It wastes the time of the user, who reads texts containing the same information several times.

Substantive problems

Substantive problems may lead to the generation of erroneous or skewed data. Due to the presence in Google Scholar search results of publications that, due to their nature, have not been subjected to a thorough review process, it is particularly important to be critical of the information obtained. An additional risk, independent of the search engine used, are errors resulting from the deliberate falsification of results or the use of artificial intelligence to create publications (Májovský et al., 2023).

Many problems are caused by frequent changes in the systematics of living organisms. Publications from different years differ in nomenclature; for example, it may occur that synonyms are treated as the names of different species. Authors cannot anticipate such changes, so readers must take this risk into account and check the names in the available registers when using the literature.

It is also problematic to refer to publications to which the reader does not have access. This is often the case with old, basic items. Most often such blockages are due to the need to preserve the rights of the author or publisher, but they make it very difficult to verify certain statements, and the suspicion arises that the author is merely quoting a source after earlier publications. Copying information secondarily after other authors without checking it in the original is a reason for perpetuating misrepresentations and errors in interpretation. A certain solution may be to use the resources of libraries and their networks, which also make old collections available electronically. Although the oldest Google Scholar records date back to 1770 (Gusenbauer, Haddaway, 2020), the representativeness of older publications in search results is low (Racki, Drabek, 2013).

A problem in scientific work, which is clearly visible when describing species, is over-generalisation, which manifests itself in attributing to an object the characteristics of the whole group to which it belongs, or treating the result of a single observation as a rule. Lack of precision, i.e. omitting data that are necessary for the correct interpretation of the information presented, also leads to errors.

Anomala dubia, due to its appearance and different colour forms, is easily confused with some other representatives of the family. Such confusion can occur if authors of non-entomological publications, with no experience in beetle species determination, refer to this species in their work. Seeking the opinion of a specialist in this area or including documentary photographs may prevent the uninformed dissemination of misinformation.

Various types of translators, which can even determine the language of the source on their own, are a very big help in using scientific literature. However, they are not perfect

and sometimes the translation obtained is illogical or even wrong. Therefore, it is safer to use specialised software and check the interpretation of the text with various examples.

The basis for using the results of other researchers is to understand the capabilities and limitations of the methodology used. Its accurate presentation is usually time-consuming and, if translators have to be used, can be difficult for the reader to interpret, especially if the text deals with complex, novel designs without reference in earlier literature. A good solution, increasingly common in scientific papers, is to supplement the text with photographs, e.g. of the observation site or experimental equipment, which makes it much easier to familiarize oneself with the methodology and scope of the research.

Most publications from the resource analysed are concerned with the occurrence of *A. dubia*, usually for the insect species register of the area. Such lists sometimes include, in addition to the location, the number of individuals observed and the date of observation. Usually it is not possible to infer from them the phenology of the species, so important e.g. in the context of climate change, because these are usually not long-term series of surveys with uniform methodology, but rather random records. They are sufficient for local checklists, but it would be advisable to collect additional information on occasion, e.g. on the location or behaviour of the insect, especially as the tools now widely available make it easier to record and archive data.

SUMMARY

Google Scholar is free, available tool for i.a. literature study. It is a constantly improving search engine, but not yet perfect. Different problems were found also in the tested search. The percentage of records with technical problems, which made the use of papers impossible or difficult, was low. However, many errors in bibliography data were revealed, the correction of which required a great deal of time. Such problems are completely absent in the articles from SCOPUS and Web of Science databases, which are perfectly prepared in this regard. Also, it is commonly believed that substantive inaccuracies and errors are removed from them during detailed editorial review, but it does not exempt readers from critical analysis of information published there.

Substantive errors are a serious problem in scientific communication. There were statements in the publications from the retrieved resource that raised suspicions about their correctness, but their verification would require deeper analysis. Many of the substantive problems may be the result of imperfections in the translators, especially in the case of texts in languages other than English.

The usefulness of Google Scholar as a data source for entomological review papers depends on many factors, including the specificity of the subject matter or its geograph-

ical scope. The results of the sample search, which was the basis of the analysis, included almost all open-access published articles from the SCOPUS and Web of Science databases corresponding to the query used. In addition, the retrieved resource included peer-reviewed scientific papers from outside these databases, which is very valuable **because there are many journals, even English-language journals, which for various reasons do not seek indexing in international prestigious bibliographic databases, but meet all the criteria for high scientific quality.** Google Scholar also lists publications of other origins, including non-peer-reviewed publications. The data from these items need to be verified in other sources, but can be the basis for interesting research topics. The retrieved records lacked much of the information needed for a comprehensive characterisation of the studied species, but this is largely the result of their absence from digitalised resources, not the limited capabilities of the search engine.

REFERENCES

Names and titles which were given in original only in Cyrillic were transliterated acc. to the norm PN-ISO 9:2000

- Aleksandrowicz O., Marczak D., Pobiedziński A., Kapuściński H., 2004.** Summer occurrence of scarab beetles (*Coleoptera, Scarabaeidae*) of the Słowiński National Park, (1997, 2002 – 2003). *Parki Narodowe i Rezerваты Przyrody*, 23: 504-511. (in Polish + summary in English)
- Amolin O.V., 2005.** An annotated list of the scoliid wasps (Hymenoptera: Scoliididae) of the Donetsk region. pp. 79-83. In: *Problems of ecology and nature protection of technogen region. The interdepartmental collection of scientific works*, no. 5, Doneck DonNU.
- Atamas' N.S., Loparev S.A., 2015.** Trofičeskie svāzi čajki hohotun'i Larus cachinnans na Srednem Dnepre. *Russkij ornitologičeskij žurnal*, 24(1176): 2846-2858. (in Russian)
- Audras G., Schaefer L., Dufay C., 1953.** Catalogue des Lucanides, Scarabéides de la région lyonnaise (suite et fin). *Bulletin mensuel de la Société linnéenne de Lyon*, 22(8): 210-216, <https://doi.org/10.3406/linly.1953.7592>.
- Baban E., 2006.** Diversitatea coleopterelor (*Coleoptera: Carabidae, Silphidae, Scarabaeidae, Cerambycidae*) din ecosistemele forestiere ale Podișului Moldovei Centrale [The diversity of coleopteras (*Coleoptera: Carabidae, Silphidae, Scarabaeidae, Cerambycidae*) from the forest ecosystems of Central Moldavian Plateau]. *Academia De Științe a Moldovei, Institutul de Zoologie*, 136 pp.
- Bajc N., 2016.** Preučevanje združljivosti entomopatogenih ogorčic (Rhabditida: Steinernematidae in Heterorhabditidae) in herbicidov v laboratorijskih razmerah [Research on compatibility of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) and herbicides under laboratory conditions]. *Magistrsko delo*, Ljubljana, 43 pp.
- Bayr U., Svae Johansen N., 2022.** Effekter av vegbelysning på insekter og deres leveområder langs europa-, riks- og fylkesvegene. *NIBIO RAPPORT*, 8(105).
- Bellmann A., 2002.** Die Trogidae, Geotrupidae, Scarabaeidae und Lucanidae (*Coleoptera*) des Weser-Ems-Gebietes. *Oldenburg, DROSERA*, pp. 109-128.
- Berglind S.-Å., Engblom E., Lingdell P.-E., 1999.** Naturligt sällsynta, hotade eller förbisedda? Nattsländorna *Semblis phalaenoides* och *S. atrata* i Sverige. [Naturally rare, threatened, or overlooked? *Semblis phalaenoides* and *S. atrata* (*Trichoptera: Phryganeidae*) in Sweden]. *Entomologisk Tidskrift*, Lund, Sweden, 120 (1-2): 1-16, ISSN 0013-886x.
- Bilokon' M.V., Zavada M.M., Cilurik A.V., 2005.** Sanitarna ocinka bukovih lisostaniv Pivničnoj Bukovini. *Naykovi dopovigi NAU*, 1(1), 13 pp.
- Bohac C., Harrison J.G., Ahrens D., 2020.** Comparative morphology of antennal surface structures in pleurostict scarab beetles (*Coleoptera*). *Zoomorphology*, 139: 327-346.
- Borowski J., Byk A., Mokrzycki T., 2016.** Beetles (*Coleoptera*) of the Rogów region. Part IX – superfamily Scarabaeoidea: Bolboceratidae, Geotrupidae, Lucanidae, Trogidae and Scarabaeidae. *World scientific news*, 44: 123-142.
- Bukejs A., 2006.** Materials about the fauna of beetles (*Insecta: Coleoptera*) of Naujene rural municipality (Daugavpils district, Latvia). Part 1. *Acta Biologica Universitatis Daugavpiliensis*, 6(1-2): 65-76.
- Burakowski B., Mroczkowski M., Stefańska J., 1983.** Chrzęszcze *Coleoptera. Scarabaeoidea, Dascilloidea, Byrrhoidea i Parnoidea*. *Katalog fauny Polski*, Nr 38, Państwowe Wydawnictwo Naukowe, Warszawa.
- Carpaneto G.M., Piattella E., Dellacasa G., Dellacasa M., Pittino R., Mazziotta A., 2011.** The lamellicorn beetles of southern Sardinia (*Coleoptera: Scarabaeoidea*). *Conservazione habitat invertebrati*, 5: 353-387.
- Chaplygina A.B., Gramma V.N., Bondarets D.I., Savynska N.O., 2015.** Arthropods in trophic-cenosis structure of collared flycatcher consortium in conditions of forest ecosystems of North-Eastern Ukraine. *Visnik Dnipropetrovs'kogo universitetu. Seriâ Biologiâ, ekologiâ*, 23(1): 74-85, doi: 10.15421/011511. (in Ukrainian + summary in English)
- Chaplygina A.B., Yuzyk D.I., Savynska N.O., 2016.** The robin, *Erithacus rubecula* (*Passeriformes, Turdidae*), as a component of heterotrophic consortia of forest cenoses, northeast Ukraine. PART 2. *Vestnik zoologii*, 50(6): 493-502.
- Chaplyhina A.B., Yuzyk D.I., Savynska N.O., Hramma V.M., 2022.** Invertebrates in the diet of collared flycatcher (*Ficedula albicollis*) nestlings in transformed forest ecosystems of north-eastern Ukraine. *Baltic Forestry*, 28(1), article id 439, 13 pp., <https://doi.org/10.46490/BF439>.
- Charrier S., 2002.** Clé de détermination des Coléoptères Lucanides et Scarabéides de Vendée et de l'Ouest de la France. *Le Naturaliste Vendéen*, 2: 61-93.
- Chenchouni H., Menasria T., Neffar S., Chafaa S., Bradai L., Chaibi R., Mekahlia M.N., Bendjoudi D., Bachir A.S., 2015.** Spatiotemporal diversity, structure and trophic guilds of insect assemblages in a semi-arid Sabkha ecosystem. *PeerJ* 3:e860; doi: 10.7717/peerj.860.
- Chittaro Y., Sanchez A., 2015.** Inventaire des coléoptères saproxyliques d'un site exceptionnel: la Châtaigneraie de Fully (VS). *Bulletin Murithienne*, 133: 13-27.
- Cosandey V., Chittaro Y., Sanchez A., 2017.** Liste commentée des Scarabaeoidea (*Coleoptera*) de Suisse. *Alpine Entomology*, 1: 57-90, doi: 10.3897/alpento.1.21179.
- De Pablo F., 2000.** Alimentación de la Lechuza Común (*Tyto alba*) en Menorca. [The diet of the barn owl (*Tyto alba*) on the isle of Minorca]. *Bolletí de la Societat d'Història Natural de les Balears*, 43: 15-26, ISSN 0212-260X.

- de Rond J., 2016.** De invloed van verstuivingsdynamiek op de insectenfauna van zeeduinen. Nulmeting en effectonderzoek van de entomofauna in de Noordwestelijke natuurkern van Nationaal Park Zuid-Kennemerland. Eindrapportage, maart 2016, 85 pp.
- Décobert O., 2010.** Notes on some European Rutelids and Cetonids. Scarabs, Occasional issue Number 54: 5-9.
- Delwaide M., Thieren Y., 2010.** Liste des coléoptères observés dans l'ancienne carrière sablonneuse de Vance (Province de Luxembourg, Belgique). Entomologie faunistique – Faunistic Entomologie, 62(1): 3-10.
- Dikmen F., Özüluğ O., 2018.** Insect (Coleoptera and Orthoptera) species of İstanbul in the Zoology Collection of İstanbul University. Turkish Journal of Bioscience and Collections, 2(1): 27-44.
- Doğan G., 2022.** Google Scholar as a data source for research assessment in the social sciences. pp. 162-180. In: Handbook on Research Assessment in the Social Sciences.
- Dumas Y., 2019.** Que savons-nous de l'Érable négondo *Acer negundo* L.? *Natura*, 2019(10): 257-283, <https://doi.org/10.5852/naturae2019a10>.
- Dutto M., 2007.** I coleotteri Scarabeoidei Pleurosticti (Coleoptera, Scarabaeoidea) e Lucanidae (Coleoptera, Lucanoidea) del Bosco del Merlino (Caramagna Piemonte, Italy). *Rivista Piemontese di Storia Naturale*, 28: 261-274
- Egorov L.V., Ručín A.B., Semišin G.B., 2017.** Materialy k poznaniū koleopterofauny Mordovskogo Gosudarstvennogo prirodnogo Zapovednika. Soobšenie 6. Trudy Mordovskogo Gosudarstvennogo prirodnogo Zapovednika imeni P.G. Smidoviča. (in Russian)
- Fătu A.-C., Dinu M.-M., Andrei A.-M., 2016.** Testing BioMel-Con bioinsecticide in some forest nurseries from Bacau county. *Romanian Journal for Plant Protection*, 9: 1-4.
- Fatu A.-C., Dinu M.M., Andrei A.M., Ciornei C., 2017.** Efficacy of biological product BIOMELCON (*Beauveria brongniartii*) during the second year of application in forest nurseries of Moldova. *Romanian Journal for Plant Protection*, X: 18-22.
- Forshage M., 2000.** Svenska namn på inhemska bladhorningar. *Entomologisk Tidskrift*, 121: 99-118.
- Frantsevich L., 2004.** Righting kinematics in beetles (Insecta: Coleoptera). *Arthropod Structure & Development*, 33: 221-235.
- Frantsevich L., Gorb S., Radchenko V., Gladun D., Polilov A., 2014.** Lehr's fields of campaniform sensilla in beetles (Coleoptera): Functional morphology. I. General part and allometry. *Arthropod Structure & Development*, 43, 523e535.
- Galinovskij N.G., Demidenko O.M., Potapov D.V., Averin V.S., 2019.** Ocenka vlianiâ vyseva zlakovyh trav v zone otčuzhdeniâ neftânyh skvažin na vidovuū strukturu soobšestv žestkokrylyh (Ectognatha, Coleoptera) na primere Sudovickogo neftânogo mestoroždeniâ (Respublika Belarus'). *Izvestiâ Gomel'skogo Gosudarstvennogo Universiteta imeni F. Skoriny*, 3(114): 11-17. (in Russian + summary in English)
- Gallou P., 1985.** Contribution à l'inventaire entomologique d'Eure-et-Loir. *Coleopteres Scarabeoidea (Lamellicornes)*, 12 pp.
- Garcia del Pino F., Palomo A., 1995.** A new strain of *Steinernema anomali* (Kozodoi, 1984) from Spain. pp. 110-111. In: COST 819 BIOTECHNOLOGY. Ecology and transmission strategies of entomopathogenic nematodes; ed.: Griffin C.T., Gwynn R.L., Masson J.P.
- Georgi R., Müller M., 2015.** Biotic risk factors in short rotation coppice in Germany: current situation, new findings and future perspectives. pp. 199-216. In: *Bioenergy from dendromass for the Sustainable Development of Rural Areas*; eds: Manning D.B., Bemann A., Bredemeier M., Lamersdorf N., Ammer Ch.
- Golawski A., 2006.** Pokarm gąsiorka *Lanius collurio* w krajo-
brazie rolniczym wschodniej Polski. *Notatki ornitologiczne*, pp. 208-213.
- Gradinarov D., Petrova J., 2011.** First report of entomopathogenic nematodes from genus *Steinernema*, 'Glaseri' group from Black Sea coast of Bulgaria. *Acta Entomologica Bulgarica*, 14(1/2): 49-55.
- Gradinarov D., Petrova E., Waeyenberge L., Karadjova O., 2011.** First report of the entomopathogenic nematode *Steinernema arenarium* (Steinernematidae: Rhabditida) in Bulgaria. *Nematologia Mediterranea*, 39: 47-52.
- Greven H., 2011.** Johann Leonhard Frisch (1666-1743) – ein wenig bekannter Pionier entomologischer Forschung. *Entomologie heute*, 23: 145-206.
- Grewal P.S., Gaugler R., Shupe Ch., 1996.** Rapid changes in thermal sensitivity of entomopathogenic nematodes in response to selection at temperature extremes. *Journal of Invertebrate Pathology*, 68: 65-73.
- Gusenbauer M., Haddaway N.R., 2020.** Which academic search systems are suitable for systematic reviews or meta-analyses? Evaluating retrieval qualities of Google Scholar, PubMed, and 26 other resources. *Research Synthesis Methods*, 11(2): 147-313, doi: 10.1002/jrsm.1378.
- Gusenleitner J., Madl M., Schedl W., Wiesbauer H., Zettel H., 2008.** Zur Kenntnis der Scoliidae (Hymenoptera) Österreichs. *Beiträge zur Entomofaunistik*, Wien, 8: 55-68.
- Gutowski J.M., Buchholz L., Kubisz D., Ossowska M., Sućko K., 2006.** Chrzęszcze saproksyliczne jako wskaźnik odkształceń ekosystemów leśnych borów sosnowych [Saproxylic beetles as indicator of deformation of pine forest ecosystems]. *Leśne Prace Badawcze*, 4: 101-144.
- Han G.Y., Lee D.W., Choo Y.M., Choo H.Y., 2012.** Effect of harvest time and infective juvenile size of the entomopathogenic nematode, *Steinernema arenarium*, on pathogenicity, development, and propagation. *Korean Journal of Applied Entomology*, 51(1): 9-18, <http://dx.doi.org/10.5656/KSAE.2012.01.1.058>. (in Korean + English summary and descriptions)
- Harčenko L.P., Čaplygina A.B., Hohlov A.N., 2002.** Pitanie drozdov v transformirovannyh landšaftah harkovskoj oblasti. *Kavkazskij ornitologičeskij vestnik*, Stavropol, 14: 99-105. (in Russian)
- Hebda G., Kata K., Żmihorski M., 2019.** The last meal: large insects predominate the diet of the European Roller *Coracias garrulus* prior to population extinction. *Bird Study*, 66(2): 173-177, doi: 10.1080/00063657.2019.1630361.
- Heijerman T., Turin H., 2017.** *Tiphia villosa*, een merkwaardige vondst van een gemutileerd exemplaar (Hymenoptera: Tiphidae). *Entomologische Berichten*, 77(1): 2-5.
- Hendrich L., Moriniere J., Haszprunar G., Hebert P.D.N., Hausmann A., Köhler F., Balke M., 2015.** A comprehensive DNA barcode database for Central European beetles with a focus on Germany: adding more than 3500 identified species to BOLD. *Molecular Ecology Resources*, 15: 795-818, <https://doi.org/10.1111/1755-0998.12354>.

- Hielkema A.J., 2022.** The referential abbreviations used by Linnaeus and other authors from 1758 through 1779 for describing species of Scarabaeoidea (Insecta: Coleoptera). *Insecta Mundi*, 0949: 1-93.
- Horvath G., Blaho M., Egri A., Hegedus R., Szel G., 2014.** Circular polarization vision of scarab beetles. pp. 147-170. In: *Polarized light and polarization vision in animal sciences*; ed. Horvath G.
- Jäger O., Brunk I., Lorenz J., 2014.** Zur Insekten- und Spinnenfauna der Kleinraschützer Heide bei Großenhain in Sachsen – Allgemeiner Teil und Käfer (Coleoptera). *Sächsische Entomologische Zeitschrift*, 8: 30-67.
- Jakubska A., Kadej M., 2006.** Preliminaries to the study of pollination ecology of *Epipactis palustris* (L.) Crantz (Orchidaceae, Neottieae) in Poland. *Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis*, III: 34-38.
- Jakubska-Busse A., Kadej M., 2008.** Pollination ecology of Marsh Helleborine *Epipactis palustris* (L.) Crantz on the Polish side of the Orlickie Mts. (the Central Sudety Mts.). *Scripra (Scripta) Facultatis Rerum Naturalium Universitatis Ostraviensis*, 186: 247-252.
- Jakubska-Busse A.J., Kadej M., 2011.** The pollination of *Epipactis* Zinn, 1757 (Orchidaceae) species in central Europe – the significance of chemical attractants, floral morphology and concomitant insects. *Acta Societatis Botanicorum Poloniae*, 80(1): 49-57.
- Jameson M.L., Paucar-Cabrera A., Solis A., 2003.** Synopsis of the New World genera of Anomalini (Coleoptera: Scarabaeidae: Rutelinae) and description of a new genus from Costa Rica and Nicaragua. *Entomological Society of America*, 6(4): 415-432.
- Jameson M.L., Mico E., Galante E., 2007.** Evolution and phylogeny of the scarab subtribe Anisopliina (Coleoptera: Scarabaeidae: Rutelinae: Anomalini). *Systematic Entomology*, 32: 429-449.
- Janiuk M., Gantner M., 2012.** Phytophagous beetles (Coleoptera) occurring on blueberry shrubs (*Vaccinium corymbosum* L.) on plantations in the environs of Lublin. *Progress In Plant Protection/Postępy w Ochronie Roślin*, 52(4): 831-836. (in Polish + summary in English)
- Japoshvili G., Hilszczański J., Byk A., Jaworski T., Łoś K., Borowski J., Tarwacki G., Piętka J., Plewa R., 2022.** New records of Coleoptera from Lagodekhi Protected Areas, with new records for Georgia (Sakartvelo). *Caucasiana*, 1: 29-39, doi : 10.3897/caucasiana.1.e85239.
- Jarry D.-T., 1964.** Les Oxyuroïdes de quelques Arthropodes dans le Midi de la France. *Mémoires originaux. Annales de Parasitologie (Paris)*, 39(4): 381-508.
- Kavka M., Veverka T., Hejkal J., 2022.** Zajímavé nálezy brouků (Coleoptera) na písčítých polích u Brambor a Horušic (střední Čechy) [Interesting findings of beetles (Coleoptera) on sandy fields near the Brambor and Horušice villages (central Bohemia). *Elateridium*, 16: 157-171.
- Klopfenstein D.V., Dampier W., 2021.** Commentary to Gusenbauer and Haddaway 2020: Evaluating retrieval qualities of Google Scholar and PubMed. *Research Synthesis Methods*, 12(2): 126-135.
- Kofler A., 1999.** Begleitfänge in Lassendorf (Kärnten) (Insecta: Coleoptera). *Carinthia II* 189/109: 617-630.
- Kolasa M., Kubisz D., Gutowski J.M., Ścibior R., Mazur M., Holecová M., Kajtoch L., 2018.** Infection by endosymbiotic “male-killing” bacteria in Coleoptera. *Folia Biologica (Kraków)*, 66(4): 165-177.
- Kolk A., Grodzki W., 2011.** Główne problemy ochrony lasu w Polsce w roku 2010 i prognoza na rok 2011. *Zpravidaj ochrany lesa 2011*, Svazek 15, Sbornik referatu z celostatniho seminare s mezinardni ucasti Skodlivi cinitele v lesich Ceska 2010/2011, Pruhonice, 2011, ISSN 1211-9342. (in Polish)
- Komaromi N.A., 2020.** Žuki pidrâdu Polyphaga (Coleoptera) gerpetobiû urbocenoziv m. Harkova. *Harkiv*, 221 pp. (in Ukrainian)
- Köteles L., Bakonyi G., 1996.** First record on the Scarabaeoidea (Coleoptera) fauna of Gödöllő (Hungary). *Folia Entomologica Hungarica*, LVII: 97-104.
- Kozel A.V., 2012.** Biologičeskoe obosnovanie meropriâtij po zaštite ot plastinčatousyh-rizofagov (*Coleoptera*, *Scarabaeidae*) sosnovyh kultur, razmešennyh na zemlâh, vyvedennyh iz selskohozâjstvenyh ugodij. *Avtoferat dissertacii na soiskanie učenoj stepeni kandidata selskohozâjstvennyh nauk po specialnosti 06.01.07 – zaštita rastenij*. (in Russian)
- Kreil F.-T., 1993.** Die Lamellicornia (Coleoptera) der Käfersammlung Dr. Theodor Hieber in den Naturkundlichen Sammlungen der Stadt Ulm, Bundesrepublik Deutschland. *Mill. d. Ver. f. Naturwiss. u. Math. Ulm (Donau)* 36/37: 49-87 1993/95.
- Kubicka A., 1981.** Scarabaeids (Coleoptera, Scarabaeidae) of Warsaw and Mazovia. *Memorabilia Zoologica*, 34: 145-164.
- Kühne S., Enzian S., Jüttersonke B., Freier B., Forster R., Rothert H., 2000.** Beschaffenheit und Funktion von Saumstrukturen in der Bundesrepublik Deutschland und ihre Berücksichtigung im Zulassungsverfahren im Hinblick auf die Schonung von Nichtzielarthropoden. In: *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft Berlin-Dahlem, Berlin, Heft 378*, 128 pp.
- Kulikov N.I., Naumkin V.P., 2004.** Plant - insect relation in buckwheat agrocoenosis. pp. 505-508. In: *Advances in Buckwheat Research. Proceedings of the 9th International Symposium on Buckwheat, Prague – Suchdol, 18-22 August 2004*.
- Kuntze R., 1936.** Krytyczny przegląd szkodników z rzędu chrząszczy zarejestrowanych w Polsce w latach 1919 - 1933. *Rocznik Ochrony Roślin*, III(2), Warszawa, 116 pp.
- Kušková K., 2014.** Brownfields jako náhradní stanoviště pro rostliny a živočichy. *Bakalářská práce, Jihočeská univerzita v Českých Budějovicích*, 39 pp.
- Lammerts E.J. Petersen J., Hochkirch A., 2009.** Beaches and Dunes. *Quality Status Report 2009, Thematic Report No. 15, WADDEN SEA ECOSYSTEM No. 25*, 20 pp.
- Lanszkia J., Giannatosb G., Heltaic M., Legakisb A., 2009.** Diet composition of golden jackals during cub-rearing season in Mediterranean marshland in Greece. *Mammalian Biology, Zeitschrift für Säugetierkunde*, 74: 72-75.
- Laznik Ž., Trdan S., 2013.** Morfološke karakteristike ogrcev s travinja, potrebne za hitro determinacijo. *Zbornik predavanj in referatov 11. Slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo Bled*, 5.–6. marec 2013, pp. 320-326.
- Laznik Ž., Trdan S., 2014.** Evaluation of different soil parameters and wild boar (*Sus scrofa* [L.]) grassland damage. *Italian Journal of Animal Science*, 13:4, 3434, doi: 10.4081/ijas.2014.3434.
- Laznik Ž., Trdan S., 2015.** Biotično zatiranje ogrcev (Coleoptera: Scarabaeidae) na travinju: izkušnje iz Gotenice. *Zbornik predavanj in referatov 12. Slovenskega posvetovanja o var-*

- stvu rastlin z mednarodno udeležbo Ptuj, 3.-4. marec 2015, pp. 307-313.
- Leontiev V.V., 2021.** The entomofauna of scarabaeoidea beetles (Coleoptera: Scarabaeoidea) in the Republic of Tatarstan. *Sarmarskij naučnyj vestnik*, 10(3): 64-71. (in Russian + summary in English)
- Leženina I.P., Gramma V.N., Savinskaja N.A., Čaplygina A.B., 2011.** Food ration of nestlings of white-collared flycatcher (*Ficedula albicollis* (Temminck, 1815)) in the plane oak forests in Kharkiv region. *The Kharkov Entomological Society Gazette*, XIX(1): 39-46. (in Russian + summary in English)
- Lila G., 2014.** Contributions to the knowledge of research on beetle parasite fauna (insects). NOTE 2. Muzeul Olteniei Craiova. Oltenia. Studii și comunicări. Științele Naturii, 30(2): 105-120, ISSN 1454-6914.
- Májovský M., Černý M., Kasal M., Komarc M., Netuka D., 2023.** Artificial intelligence can generate fraudulent but authentic-looking scientific medical articles: Pandora's box has been opened. *Journal of Medical Internet Research*, 25, e46924.
- Maksimova V.I., 2008.** Kak otličít' ličínok âponskogo žuka ot nekarantinnyh vidov plastinčatousyh žukov. *Zašita i karantin rastenij*, Karantin, 5: 41-43. (in Russian)
- Malicky H., 1975.** Vergleich der Scarabaeiden-Faunen (Coleoptera) Mitteleuropas (s.l.) und Jugoslawiens. pp. 329-341. In: *Verhandlungen des Sechsten Internationalen Symposiums über Entomofaunistik in Mitteleuropa, Lunz am See (Österreich)*, 1.-6. September 1975.
- Marczak D., Mroczyński R., 2016.** A contribution to the knowledge of the fauna of the Kampinos National Park: Scarabaeidae. Part 1. Subfamilies: Melolonthinae, Sericinae, Rutelinae, Dynastinae i Cetoninae. *Entomological News*, 35(3): 161-171.
- Maréchal P., 1993.** Over externe factoren die de habitatkwaliteit van de Grauwe Klauwier *Lanius collurio* beïnvloeden. *Hel Vogeljaar*, 41: 34-48.
- Martikainen P., 2016.** Hailuodon Mäntyniemen ja Huilunnon, Siikajoen Tavon ja Uudenkaarlepyyn Storsandenin kovakuoriaiskartoitukset 2015. *Paahde Life (LIFE13NAT/FI/000099)*.
- Meleshko J.Y., 2003.** The species composition of the beetles (*Coleoptera*) on the poplars in Minsk. *Second International Symposium on Plant Health in Urban Horticulture, Berlin, Germany, August 27-29, 2003, Heft 394*: 58-60.
- Mertlik J., 2011.** Příspěvek k ochraně písčitých biotopů východních Čech s uvedením nálezu devíti vzácných psamofilních brouků [Contribution to the conservation of sand biotopes of the Eastern Bohemia with citation of findings of nine rare psammophilous beetles]. *Elateridarium*, 5: 5-42.
- Mertlik J., Brůha P., 2011.** Výskyt *Dicronychus equiseti* (Coleoptera: Elateridae) v Čechách (Česká republika) [The occurrence of *Dicronychus equiseti* (Coleoptera: Elateridae) in Bohemia (Czech Republic)]. *Elateridarium*, 5: 248-253.
- Mikheyev A.V., 2002.** Comparative characteristics of genus *Martes* diet in forest ecosystems of Ukraine steppe zone. *Vestnik zoologii*, 36(3): 45-54. (in Russian).
- Morelli F., Bussièrec R., Goławski A., Tryjanowski P., Yosef R., 2015.** Saving the best for last: Differential usage of impaled prey by red-backed shrike (*Lanius collurio*) during the breeding season. *Behavioural Processes*, 119: 6-13, <http://dx.doi.org/10.1016/j.beproc.2015.07.006>.
- Nadvorný V.G., 2007.** Biodiversity and biotope distribution of invertebrates on the right side of the dam in Kremenchuk Reservoir (Cherkassy Suburb). *ZOOCENOSIS*, pp. 30-32 (in Ukrainian).
- Naumkin V.P., Lysenko N.N., 2014.** Buckwheat in the conservation of insects species diversity. *Vestnik OrelGAU*, 4(49): 12-18.
- Negrobov S.O., 2009.** Seasonal dynamics of scarab beetles (Coleoptera, Lamellicornia) in Voronezh Province. *Entomological Review*, 89(1): 113-115.
- Netoiu C., Tomescu R., Badela O., Nica M.S., 2007.** Infestation and damage characteristics produced by some chafers species in young plantations established on sandy soils from the south of Oltenia region. *Analele ICAS*, 50: 239-259.
- Nguyen K.B., Smart G.C. jr., 1993.** Scanning electron microscope studies of *Steinernema anomali* Kozodoi, 1984. *Journal of Nematology*, 25(3): 486-492.
- Nicolas J.-P., Nicolas J.-L., 1965.** Révision commentée de la collection et du catalogue des coléoptères de la Côte-d'Or d'A. Rouget (1er note). *Bulletin mensuel de la Société linnéenne de Lyon*, 34^e année, n°1 : 10-22, <https://doi.org/10.3406/linly.1965.5725>.
- Nijssen M., Geertsma M., 2014.** Het dieet van Limburgse Grauwe Klauwieren. *Limburgse Vogels*, 24: 45-54.
- Niklas O.F., 1963.** Infektionen durch *Rickettsiella melolonthae* (Krieg) Philip bei larven von *Amphimallon solstitiale* (Linnaeus), *Anomala dubia aenea* (De Geer), *A. (Phyllopertha) horticola* (Linnaeus), *Maladera (Serica) brunnea* (Linnaeus) und *Onthophagus spec (Coleoptera: Lamellicornia)*. *Beiträge zur Entomologie*, 13(3/4): 395-401.
- Nuhlíčková S., Křištín A., Degma P., Hoi H., 2016.** Variability in nestling diet of European hoopoes: annual and sampling effect. *Folia Zoologica*, 65(3): 189-199.
- Ødegaard F., Hanssen O., Aagaard K., Aarrestad P.A., 2001.** Grunnlag for standardisert klassifisering av habitatyper og trusselfaktorer i den nasjonale rødlista. *Test av systemet på tre taksonomiske grupper*. NINA Fagrapport, 47: 1-45.
- Olsson F., 2006.** Holocene forest history at Stavsåkra, Småland, south central Sweden inferred from insect fossils. *The ESS Bulletin*, 4(1): 44-61.
- Olsson F., Lemdahl G., 2009.** A continuous Holocene beetle record from the site Stavsåkra, southern Sweden: implications for the last 10 600 years of forest and land use history. *Journal of Quaternary Science*, 24(6): 612-626, doi: 10.1002/jqs.1242.
- Orłowski G., Karg J., Czarnecka J., 2011.** Frugivory and size variation of animal prey in Black Redstart *Phoenicurus ochruros* during summer and autumn in south-western Poland. *Ornis Fennica*, 88: 161-171.
- Ostrovskij A.M., 2015.** Scarabaeoidea beetle complex (Coleoptera, Scarabaeoidea) south-east of Belarus. pp. 90-93. In: *Sbornik materialov IV Meždunarodnoj konferencii 26–28 oktábrá 2015 g., g. Tomsk, Rossiá*. (in Russian + summary in English)
- Pacheco T.L., Bohacz C., Ballerio A., Schoolmeesters P., Ahrens D., 2022.** Revisiting trends in morphology of antennal sensilla in scarabaeoid beetles. *Zoomorphology*, 141: 315-326.

- Pandey A.K., 2013.** Extent of natural infection of entomopathogens in whitegrubs of Western Himalayan region of Uttarakhand. *Journal of Biological Control*, 27(4): 340-342.
- Parra A.E.G., 2020.** Producción semisólida y almacenamiento del nemátodo entomopatógeno *Heterorhabditis bacteriophora*: Revisión de Literatura. Proyecto especial de graduación presentado como requisito parcial para optar al título de Ingeniero Agrónomo en el Grado Académico de Licenciatura. Zamorano, Honduras, 28 pp.
- Pedersen L., Geertsma M., Tøttrup A.P., 2012.** Prey diversity is affected by climate and differs between age classes in the Red-backed Shrike (*Lanius collurio*). *Ornis Fennica*, 89: 99-108.
- Pereira F.A., Mugnaini R., 2023.** Mapping the use of Google Scholar in evaluative bibliometric or scientometric studies: A bibliometric review. *Quantitative Science Studies*, 4(1): 233-245.
- Peters A., Galarza M., 2004.** Efficacy of long *Steinernema* spp. nematodes against scarabs. *Environmental Science, Biology*.
- Petrova Y., Gradinarov D., Chehlarov E., Teofilova T., Kodzhabashev N., 2022.** Scarabaeoid beetles (Coleoptera: Scarabaeoidea) in Sarnena Sredna Gora Mountains. In: *Fauna of Sarnena Sredna Gora Mts*, eds: Georgiev D., Bechev D. Yancheva V., Part 3 ZooNotes, Supplement 11.
- PIORIN, 2016.** *Popillia* japońska (*Popillia japonica* Newmann). Szkodnik podlegający obowiązkowi zwalczania. https://piorin.gov.pl/files/userfiles/giorin/publikacje/ulotki/p_japonica_wyd.1.pdf, 2 pp.
- Poinar Jr. G.O., Kozodoi E.M., 1988.** *Neoaplectana glaseri* and *N. anomali*: sibling species or parallelism? *Revue Nématol.*, 11(1): 13-19.
- Pokarzhevskii A., Zhulidov A., 1995.** Halogenes in soil animal bodies: A background level. In: *Contaminated Soil; '95. Soil & Environment*, eds.: Van Den Brink W.J., Bosman R., Arendt F., vol 5. Springer, Dordrecht, https://doi.org/10.1007/978-94-011-0415-9_103.
- Polat A., Yildirim E., Uliana M., 2018.** A contribution to the knowledge of the Dynastinae, Rutelinae and Melolonthinae fauna of Turkey (Coleoptera: Scarabaeidae). *Entomofauna*, 39/2, 08: 597-614.
- Prakhodskiy S.A., Kaplich V.M., Jarmolovich V.A., 2010.** Noxious organisms of scots pine (*Pinus sylvestris* L.) in forest plantations. pp. 491-497.
- Prokof'eva I.V., 2003.** K pitaniû ptencov derâby *Turdus viscivorus* i černogo drozda *Turdus merula* na ûge Leningradskoj oblasti. *Russkij ornitologičeskij žurnal*, 219: 423-424. (first publication in 1983 r.) (in Russian)
- Prokof'eva I.V., 2007.** Sravnenie pitaniâ drozdov-derâb *Turdus viscivorus* v raznyh mestnostâh, silno udalennyh drug ot druga. *Russkij ornitologičeskij žurnal*, Tom 16, Èkspress-vypusk 340: 45-49. (in Russian)
- Prokof'eva I.V., 2013.** Pitanie lugovyh vorob'inyh ptic v period gnezdovaniâ. *Russkij ornitologičeskij žurnal*, Tom 22, Èkspress-vypusk, 895: 1817-1822. (in Russian)
- Quintero Marin M.P., 2003.** Comparacion en laboratorio de la patogenicidad de tres especies nativas de nematodos entomopatógenos (Rhabditida) sobre larvas de tercer instar de *Phyllophaga menetriesi* (Blanchard) (Coleoptera: Scarabaeidae). Trabajo de grado presentado como requisito parcial para optar al título de Bióloga, 49 pp.
- Racki G., Drabek A., 2013.** Cytowania i wskaźnik Hirscha: gdzie szukać, jak obliczać? *Forum Akademickie*, 02/2013, <https://prenumeruj.forumakademickie.pl/fa/2013/02/cytowania-i-wskaznik-hirscha-gdzie-szukac-jak-obliczac/>.
- Radics L., Mikóházi D., 2010.** Principles of common buckwheat production. *The European Journal of Plant Science and Biotechnology*, pp. 57-63.
- Radzikowski P., Matyka M., Berbeć A.K., 2020.** Biodiversity of weeds and arthropods in five different perennial industrial crops in eastern Poland. *Agriculture*, 10, 636; doi: 10.3390/agriculture10120636.
- Ramírez-Ponce A., Morón M.A., 2009.** Relaciones filogenéticas del género *Anomala* (Coleoptera: Melolonthidae: Rutelinae) [Phylogenetic relationships of the genus *Anomala* (Coleoptera: Melolonthidae: Rutelinae)]. *Revista Mexicana de Biodiversidad*, 80: 357-394.
- Reiter W., 1986.** Phänologie und Dominanzwechsel phytophager Insekten am Sanddorn (*Hippophaë rhamnoides* L.) anhand von Untersuchungen in Nord- und Südtirol. *Berichte des naturwissenschaftlichen-medizinischen Verein, Innsbruck*, 73: 169-186.
- Reitze M., Nentwig W., 1991.** Comparative investigations into the feeding ecology of six Mantodea species. *Oecologia*, 86: 568-574.
- Romero-Lopez A.A., Arzuffi R., Valdez J., Sanchez-Espindola E., Moro M.A., 2011.** Tissues involved in sex pheromone production in *Phyllophaga obsoleta* (Coleoptera: Scarabaeoidea: Melolonthidae). *Annals of the Entomological Society of America*, 104(5): 960-965, doi: 10.1603/AN10068.
- Ruta R., Orzechowski R., Aleksandrowicz O., Borowski J., Buchholz L., Komosiński K., Lubecki K., Przewoźny M., 2016.** Chrząszcze (Insecta: Coleoptera) Gryżyńskiego Parku Krajobrazowego [Beetles (Insecta: Coleoptera) of Gryżyński Landscape Park], *Przegląd Przyrodniczy*, XXVII(2): 28-62.
- Sauvayre R., 2022.** Types of errors hiding in Google Scholar data. *Journal of Medical Internet Research*, 24(5): e28354, doi: 10.2196/28354.
- Sažnev A.S., Anikin V.V., 2021.** Žestokrylye nadsemejstva Scarabaeoidea (COLEOPTERA) Nacional'nogo Parka «Hvalynskij» (Saratovskaâ oblast'). *Èntomologičeskije i Parazitologičeskije Issledovaniâ v Povolž'e*, 18: 20-26. (in Russian)
- Schacht W., 2017.** Zur Bedeutung von Kiesgruben für den Erhalt der Artenvielfalt am Beispiel der Käfer einer Elbe-nahen Grube im nordöstlichen Niedersachsen (Coleoptera). *Entomologische Zeitschrift Schwanfeld*, 127(1): 17-28.
- Schacht W., 2022.** Zum 275. Geburtstag des Koleopterologen Franz von Paula Schrank (Coleoptera). *Entomologische Zeitschrift, Schwanfeld*, 132(3): 153-156.
- Scharfy J., 2012.** Die Entwicklung der Dolchwespe *Scolia hirta* (Schrank, 1781) (Hymenoptera, Scoliidae) in einem Komposthaufen in Österreich. *Ampulex*, 4: 47-50.
- Singh V.K., Srichandan S.S., Piryani R., Kanaujia A., Bhat-tacharya S., 2023.** Google Scholar as a pointer to open full-text sources of research articles: A useful tool for researchers in regions with poor access to scientific literature. *African Journal of Science, Technology, Innovation and Development*, 15(4): 450-457.
- Smith A.B.T., 2009.** Checklist and Nomenclatural Authority File of the Scarabaeoidea of the Nearctic Realm including Canada, the continental United States, and the northern Mexican

- states of Baja California, Baja California Sur, Chihuahua, Coahuila de Zaragoza, Durango, Nuevo Leon, Sinaloa, Sonora, Tamaulipas, and Zacatecas. Canadian Museum of Nature, 97 pp.
- Šohin I.V., 2011.** Analiz ekologo-geografičeskikh osobennostej fauny plastinčatousyh žukov (Coleoptera: Scarabaeoidea) ũžnoj Rossii. *Èkologiã životnyh*, 4: 160-172. (in Russian)
- Sörensson M., 2020.** Återinventering av insekter på Bititten och Palmers väg i Åhus 2020. Biosfärkontoret Kristianstads Vattenrike, Vattenriket i fokus 2020:06.
- Spaar D., Kleinhempel H., Fritzsche R., 1987.** Diagnose von Krankheiten und Beschädigungen an Kultupflanzen. Kartoffeln. Springer-Verlag, Berlin Heidelberg New York London Paris Tokyo, 136 pp., ISBN-13: 978-3-642-70916-6.
- Spaar D., Kleinhempel H., Fritzsche R., 1988.** Diagnose von Krankheiten und Beschädigungen an Kultupflanzen. Zucker- und Futterrüben. Springer-Verlag, Berlin Heidelberg New York London Paris Tokyo, ISBN-13: 978-3-642-72800-6.
- Szafraniec S., Kubisz D., Melke A., 2021.** Przegląd chrząszczy (Coleoptera) Masywu Babiej Góry [Review of beetles (Coleoptera) of Mt. Babia Góra]. pp. 163-212. In: Fauna masywu Babiej Góry. Bezkręgowce. [Fauna of Babia Góra massif. Invertebrates]
- Telfer M.G., Eversham B.C., 1995.** Invertebrate Recording on Suffolk Breckland Sites of Special Scientific Interest during 1993 and 1994. 40 pp.
- Teodorescu I., Maican S., 2014.** Diversity of entomofauna (Hymenoptera, Diptera, Coleoptera) in the Comana Natural Park (Romania). Bucharest, Romanian Journal of Biology – Zoology, 59(1): 17-34.
- Teppner H., 2008.** *Scolia hirta* (Hymenoptera-Scoliidae) neu für die Steiermark. Mitteilungen des naturwissenschaftlichen Vereines für Steiermark, Graz, 138 : 5-8.
- Théodoridès J., 1963.** Faune terrestre et d'eau douce des Pyrénées-Orientales Fascicule 8 Sporozoaires et Cnidosporidies. HERMANN, 37 pp.
- Thomaes A., Drumont A., Eylenbosch S., Raemdonck H., Muls D., Dekuijper C. Dahan L., 2016.** Three new localities for *Gnorimus nobilis* in northern Belgium (Coleoptera: Cetoniidae). Bulletin de la Société royale belge d'Entomologie/ Bulletin van de Koninklijke Belgische Vereniging voor Entomologie, 152: 122-127.
- Thomaes A., Scheers K., Van Keer K., 2022.** Establishment after transport adds *Mimela junii* (Duftschmid, 1805) to the Belgian fauna (Coleoptera: Rutelidae). Bulletin de la Société royale belge d'Entomologie / Bulletin van de Koninklijke Belgische Vereniging voor Entomologie, 158: 60-67.
- Toth M., Leal W.S., Szarukan I., Lesznyak M., Szocs G., 1994.** 2-(E)-Nonen-1-ol: male attractant for chafers *Anomala vitis* FABR. and *A. dubia* SCOP. (Coleoptera: Scarabaeidae). Journal of Chemical Ecology, 20(10): 2481-2487.
- Tóth M., Szarukán I., Furlan L., Sivcev I., Ilovai Z., Ujváry I., Szics G., 1997.** Hazai fejlesztésű szexferomon csapdák bogárkártevőkre (Poszter) [Hungarian developed sexpheromon traps for beetle (Coleoptera) pests (poster)]. 1st International Plant Protection Symposium at DAU, 19-22 August 1997, pp. 105-106.
- Tryjanowski P., Karg M.K., Karg J., 2003.** Diet composition and prey choice by the red-backed shrike *Lanius collurio* in western Poland. Belgian Journal of Zoology, 133(2): 157-162.
- Uhan T.S., 2005.** Bioefikasi nematoda entomopatogen *Steinernema* spp. isolat Lembang terhadap larva *Crocidolomia pavonana* (F) pada tanaman kubis di rumahkaca. Jurnal Hortikultura, 15(2): 109-115.
- Ungar O.J., Muhanna N., Chaushu H., 2023.** Google Scholar as a sole literature source for meta analyses in otolaryngology head and neck surgery. Indian Journal of Otolaryngology and Head & Neck Surgery, 75: 864-870, <https://doi.org/10.1007/s12070-023-03532-8>.
- Urban P., Winzer K., Schulze W., 2020.** Zum Auftreten von *Mimela junii* (DUFTSCHMID, 1805) in Nordrhein-Westfalen (Coleoptera, Scarabaeidae, Rutelinae). Bielefeld, Mitt. ArbGem. westfäl. Entomol., 36(1): 7-9.
- van der Kruis B., 2008.** Is de Grauwe Klauwier bezig met een voorzichtige opmars in de Nederlandse duinen? Fitis, 44(1): 18-21.
- van Duinen G.-J., van Kleef H., Nijssen M., van Turnhout Ch., Verberk W., Holtland J., Esselink H., 2004.** Schaal en intensiteit van herstelmaatregelen: hoe reageert de fauna? pp. 189-240. In: Duurzaam natuurherstel voor behoud van biodiversiteit. 15 jaar herstelmaatregelen in het kader van het overlevingsplan b; ed.: Van Duinen G., Bobbink R., Van Dam C.
- van Duinen G.J., Beusink P., Nijssen M., Esselink H., 2005.** Larval development of *Anomala dubia* (Scarabaeidae) in coastal dunes: Effects of sand-spray and *Ammophila arenaria* rootbiomass. Proceedings of the Netherlands Entomological Society Meeting, 16: 63-70.
- van Oosten H., van Turnhout Ch., Beusink P., Majoor F., Hendriks K., Geertsma M., van den Burg A., Esselink H., 2008.** Broed- en voedselécologie van Tapuit: Opstap naar herstel van de faunadiversiteit in de Nederlandse kustduinen. 34 pp.
- van Oosten H.H., van den Burg A.B., 2015.** Limitaties voor insectivore vogels in het duinvoedselweb. Vooronderzoek 2015, 42 pp.
- van Oosten H.H., van den Burg A.B., Versluijs R., Siepel H., 2014.** Habitat selection of brood-rearing Northern Wheat-eaters *Oenanthe oenanthe* and their invertebrate prey. Ardea, 102(1): 61-69.
- Veremeev V.N., 2005.** Èkologičeskaâ model' izmenenij kompleksov počvennoj mezofauny lesnyh èkosistem v usloviãh nedostatka vlagi. Biosystems Diversity, 13(2): 38-43. (in Russian)
- Veremeev V.N., Sinenok N.L., Černčikova O.V., 2001.** Èkologičeskij monitoring sostoãniã kompleksov počvennoj mezofauny lugov pojmy reki Sož. Izvestiã Gomelskogo gosudarstvennogo Universiteta imeni F.Skoriny, 4 Biologiã, pp. 15-19. (in Russian)
- Voigt E., Tóth M., 2004.** Three years of mass trapping with sex attractant traps for control of *Anomala* Scarabs in ripening peaches. Integrated plant protection in stone fruit. IOBC/wprs Bulletin, 27(5): 67-71.
- Voigt E., Tóth M., 2009.** Opportunities and limitations of mass trapping of difficult-to-control fruit pests: efforts in Hungary. Pheromones and other Semiochemicals. IOBC/wprs Bulletin, 41: 61-66.
- Von Klausnitzer B., Franke R., Liebig W.-H., Scholz A., 2013.** *Scolia hirta* (Schrank, 1781) (Hymenoptera, Scoliidae) und ihre Wirte (Coleoptera, Scarabaeidae) in der Oberlausitz. Berichte der Naturforschenden Gesellschaft der Oberlausitz, Görlitz, 21: 95-102.

- Von Niklas O.F., 1964.** Vertikalbewegungen Rickettsiose-krank-er Larven von *Amphimallon solstitiale* (LINNAEUS), *Anomala dubia aenea* (DE GEER) and *Maladera brunnea* (LINNAEUS) (*Col.*, *Lamelleornia*). *Anzeiger für Schädling-skunde*, 37: 22-24.
- Watkins C., Mann D., 2018a.** The status and distribution of the scarab beetles *Rhysothorax rufa* and *Onthophagus nuchicornis* on Welsh dunes in 2017. NRW Evidence Report No. 263, 40 pp.
- Watkins C.M., Mann D.J., 2018b.** The status and distribution of the scarab beetles *Rhysothorax rufa* and *Ammoecius brevis* on Lancashire dunes in 2017. 34 pp.
- What are the advantages and disadvantages of using Google Scholar for academic research? <https://www.linkedin.com/advice/0/what-advantages-disadvantages-using-google-scholar> (accessed 13.12.2023).
- Wolender M., Zych A., 2007.** Beetles (Coleoptera) from sea-side beach and dunes in the regions of Świnoujście, Międzyzdroje and Wiselka (Poland) located along the southern coast of the Baltic Sea. *Baltic Journal of Coleopterology*, 7(1): 61-71.
- Wouters B., van Oosten H., 2013.** Kink in de (voedsel)keten van duingraslanden? *De Levende Natuur*, 114(5): 232-237.
- Wu S., 2013.** Efficacy of entomopathogenic nematodes and entomopathogenic fungi against masked chafer white grubs, *Cyclocephala* spp. (Coleoptera: Scarabaeidae). Dissertation for the degree of Doctor of Philosophy in Entomology, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 135 pp.
- Zhukov A.V., 2006.** Soil animals gygromorphs and their diagnostic importance for gygrotops indicating. 17 pp., https://www.zoology.dp.ua/wp-content/downloads/zhukov/ZH_06_02.pdf. (in Russian + summary in English)
- Zhukov A.V., 2007.** Soil animal trophocoenomorphes and their diagnostic importance for gygrotops indicating. 22 pp., https://www.researchgate.net/publication/273379736_TROFO-CENOMORFY_POCVENNYH_ZIVOTNYH_I_IH_DIAG-NOSTICESKOE_ZNACENIE_DLA_USTANOVLENIA_TROFOTOPOV. (in Russian + summary in English)
- Zúber M., 2011.** Listorozi brouci (Coleoptera: Scarabaeoidea) dolního Pojizeří [Scarabaeoidea of the lower Jizera river region]. *Elateridarium*, 5: 43-54.

Author

ORCID

Alina Bochniarz

0000-0001-6545-3041

received – 14 December 2023

revised – 19 December 2023

accepted – 27 December 2023

Author declares no conflict of interest.



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike (CC BY-SA) license (<http://creativecommons.org/licenses/by-sa/4.0/>).